

# Increase of mitochondria and mitochondrial DNA in response to oxidative stress in human cells

Hsin-Chen LEE\*, Pen-Hui YIN†, Ching-You LU\*, Chin-Wen CHI† and Yau-Huei WEI\*<sup>1</sup>

\*Department of Biochemistry and Center for Cellular and Molecular Biology, National Yang-Ming University, Taipei, Taiwan 112, Republic of China, and †Institute of Pharmacology, National Yang-Ming University and Department of Medical Research and Education, Taipei Veterans General Hospital, Taipei, Taiwan 112, Republic of China

Mitochondrial respiratory function is impaired in the target tissues of patients with mitochondrial diseases and declines with age in various human tissues. It is generally accepted that respiratory-chain defects result in enhanced production of reactive oxygen species and free radicals in mitochondria. Recently, we have demonstrated that the copy number of mitochondrial DNA (mtDNA) is increased in the lung tissues of elderly human subjects. The mtDNA copy number was suggested to be increased by a feedback mechanism that compensates for defects in mitochondria harbouring mutated mtDNA and a defective respiratory system. However, the detailed mechanism remains unclear. In this study, we treated a human lung fibroblast cell line, MRC-5, with H<sub>2</sub>O<sub>2</sub> at concentrations of 90–360  $\mu$ M. After the treatment for 24–72 h, we found that cells were arrested at G<sub>0</sub> and G<sub>1</sub> phases but that mitochondrial mass and mtDNA content were significantly increased in a concentration- and time-de-

pendent manner. Moreover, the oxidative stress induced by buthionine sulfoximine was also found to cause an increase in mitochondrial mass of the treated cells. Increased uptake of a vital mitochondrial dye Rhodamine 123 and enhanced tetrazolium [MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide] reduction revealed that the mitochondria increased by H<sub>2</sub>O<sub>2</sub> treatment were functional. In addition, the increase in the mitochondrial mass was also observed in cell-cycle-arrested cells induced by mimosine, lovastatin and genistein. Taken together, these findings suggest that the increase in mitochondrial mass and mtDNA content are the early molecular events of human cells in response to endogenous or exogenous oxidative stress through cell-cycle arrest.

**Key words:** aging, cell-cycle arrest, hydrogen peroxide, mtDNA copy number, reactive oxygen species.

## INTRODUCTION

Mitochondria are the intracellular organelles responsible for ATP synthesis through the coupling of oxidative phosphorylation to respiration in human and animal cells. Under normal physiological conditions, a small fraction of oxygen consumed by mitochondria is converted to superoxide anions, H<sub>2</sub>O<sub>2</sub> and other reactive oxygen species (ROS) [1]. Moreover, mitochondrial respiratory function has been demonstrated to decline in various human tissues during the aging process [2,3]. This is thought to be caused, at least partly, by oxidative damage and mutation of mitochondrial DNA (mtDNA) in somatic tissues of aged individuals [3].

Recently, we found that mtDNA copy number is increased in the lung tissues of elderly human subjects [4]. Moreover, we showed that mtDNA mutation, oxidative DNA damage and lipid peroxidation are increased during aging in the human lung [4–7] and other tissues [8,9]. These findings are consistent with a previous report that mtDNA contents are increased in the tissues of old rats [10] and of elderly subjects [11]. This increase in the content of mtDNA was thought to compensate for respiratory-function decline during the aging process [12]. However, the molecular mechanism underlying the increase of mtDNA associated with respiratory-function decline in human aging remains unclear.

Mitochondria are the major intracellular source and primary target of ROS, which are generated under normal conditions as by-products of aerobic metabolism in animal and human cells. It

has been established that defects in the respiratory chain lead to enhanced production of ROS and free radicals in mitochondria [13–15]. Moreover, H<sub>2</sub>O<sub>2</sub> has been proposed to play a role as a secondary messenger in signal-transduction pathways [16,17]. The increase of ROS and enhanced oxidative stress stimulates the expression of early growth-related genes such as *c-fos* and *c-jun* [18]. Recently, H<sub>2</sub>O<sub>2</sub> was proposed to be involved in the communication between mitochondria and the nucleus [19,20]. Therefore, we hypothesized that the age-dependent increase in the production of superoxide anions and H<sub>2</sub>O<sub>2</sub> from mitochondria is one of the factors involved in feedback compensation for the decline of respiratory function during the aging process [21].

To test this hypothesis, we exposed MRC-5, a human lung fibroblast cell line, to oxidative stress by treatment with low concentrations of H<sub>2</sub>O<sub>2</sub> or buthionine sulfoximine (BSO). The effect of oxidative-stress treatment on the copy number of mtDNA and mitochondrial mass in the cell was investigated.


## MATERIALS AND METHODS

### Cell culture

Human lung fibroblasts, from the MRC-5 cell line, were cultured at 37 °C in humidified 5% CO<sub>2</sub>/95% air in Dulbecco's modified Eagle's medium (Gibco/BRL, Bethesda, MD, U.S.A.) supplemented with 10% fetal bovine serum, 50 units/ml penicillin G

Abbreviations used: mtDNA, mitochondrial DNA; ROS, reactive oxygen species; BSO, buthionine sulfoximine; NAO, 10-*n*-Nonyl-Acridine Orange; DCF, 2',7'-dichlorofluorescein; DCFH-DA, 2',7'-dichlorofluorescein diacetate; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide; DiOC<sub>6</sub>(3), 3,3'-dihexyloxycarbocyanine iodide.

<sup>1</sup> To whom correspondence should be addressed (e-mail joeman@mailsv.ym.edu.tw).



**Figure 1** Cell growth and viability of MRC-5 cells treated with H<sub>2</sub>O<sub>2</sub>

MRC-5 cells were treated with H<sub>2</sub>O<sub>2</sub> at concentrations from 90 to 360 μM. After treatment for 24–72 h, cells were enumerated using a haemocytometer (A), and the number of viable cells was determined by Trypan Blue exclusion (B). Data points represent measured values ± S.D. of two independent experiments performed on cells from three separate dishes. The differences in growth kinetics and viability between untreated and H<sub>2</sub>O<sub>2</sub>-treated cells were found to be statistically significant by using one-way ANOVA (\*,  $P < 0.05$ ).

and 50 mg/ml streptomycin sulphate. Cells were subjected to a 1:2 split every 4 days.

### Oxidative-stress treatment

Oxidative-stress treatment was performed on cells that had all been cultured at the same density. An aliquot of  $2.5 \times 10^5$  cells at 25–35 passages was plated in triplicate in 25-cm<sup>2</sup> flasks 24 h before treatment. For H<sub>2</sub>O<sub>2</sub> treatment, a suitable aliquot of 30 % H<sub>2</sub>O<sub>2</sub> solution was freshly diluted into the culture medium before the experiment. Cells were exposed to the indicated concentrations of H<sub>2</sub>O<sub>2</sub> for a fixed period of time under normal culture conditions as described above. To deplete intracellular glutathione, cells were treated with 1 or 2 mM BSO (Sigma, St. Louis, MO, U.S.A.) for 72 h. The treated cells were then washed once with Hanks' balanced salt solution (Gibco/BRL) and were harvested by trypsinization at various time points for further analysis.

### Treatment with drugs affecting the cell cycle

To arrest the cell cycle, cells were treated with 200 μM mimosine, 2.5 μM lovastatin or 50 μM genistein (Sigma) for 24 h. The

treated cells were then washed once with Hanks' balanced salt solution and were harvested by trypsinization at various time points for cell-cycle analysis and mitochondrial-mass determination.

### Cell growth and viability assay

Cells were harvested by trypsinization and resuspended in 1 ml of Medium 199 (Gibco/BRL). The cells were enumerated using a haemocytometer, and the number of viable cells was determined on the basis of their exclusion of 0.4 % Trypan Blue.

### Cell-cycle analysis

After treatment with H<sub>2</sub>O<sub>2</sub> or other chemicals, cells were trypsinized and resuspended in 70 % ethanol. The cells were incubated on ice for at least 1 h and resuspended in 1 ml of cell-cycle assay buffer (0.38 mM sodium citrate, 0.5 mg/ml RNase A and 0.01 mg/ml propidium iodide) at a concentration of  $5 \times 10^5$  cells/ml. Samples were stored in the dark at 4 °C until cell-cycle analysis, which was carried out by use of a flow cytometer and ModFit LT 2.0 software (Verity Software, Topsham, ME, U.S.A.).

### Determination of mitochondrial mass

The fluorescent dye 10-*n*-Nonyl-Acridine Orange (NAO; Molecular Probes, Eugene, OR, U.S.A.), which binds specifically to cardiolipin at the inner mitochondrial membrane independently of membrane potential ( $\Delta\Psi_m$ ) [22,23], was used to monitor the mitochondrial mass [24]. Cells were trypsinized and resuspended in 0.5 ml of Medium 199 containing 10 μM NAO or 10 μM MitoTracker Green FM (Molecular Probes), which is preferentially accumulated in mitochondria regardless of mitochondrial membrane potential. After incubation for 10 min at 25 °C in the dark, cells were transferred immediately to a tube on ice for flow-cytometric analysis.

### Dichlorofluorescein staining


H<sub>2</sub>O<sub>2</sub> production by cultured cells was measured using the probe 2',7'-dichlorofluorescein diacetate (DCFH-DA; Molecular Probes) [24–26]. This probe is accumulated by cells and hydrolysed by cytoplasmic esterases to become 2',7'-dichlorofluorescein, which then reacts with H<sub>2</sub>O<sub>2</sub> to give the fluorescent product 2',7'-dichlorofluorescein (DCF) [25]. Cells were incubated with 5 μM DCFH-DA in culture medium for 30 min at 37 °C and then washed, resuspended in 0.5 ml of PBS, and submitted to flow-cytometric analysis.

### Measurement of mitochondrial membrane potential

Cells were trypsinized and resuspended in 0.5 ml of Medium 199 containing 0.5 μg/ml Rhodamine 123 (Molecular Probes) or 3,3'-dihexyloxycarbocyanine iodide [DiOC<sub>6</sub>(3); Molecular Probes]. After incubation for 30 min at 37 °C in the dark, cells were transferred immediately to a tube on ice for analysis of the fluorescence intensity by flow cytometry.

### Flow-cytometric analysis

A FACScan flow cytometer (Becton Dickinson, Bedford, MA, U.S.A.) equipped with a 488-nm argon laser was used for the flow-cytometric analysis. Forward and side scatters were used to establish size gates and exclude cellular debris from the analysis.




**Figure 2** Cell-cycle arrest induced by H<sub>2</sub>O<sub>2</sub>

MRC-5 cells were treated with 90–360 μM H<sub>2</sub>O<sub>2</sub>. After treatment for 24 h, the percentages of cells in G<sub>0</sub>-G<sub>1</sub> phase (**A**) and S phase (**B**) were analysed on a flow cytometer as described in the Materials and methods section. C, control.

The excitation wavelength was set at 488 nm. The observation wavelength of 530 nm was chosen for green fluorescence and 585 nm for red fluorescence and the intensities of emitted fluorescence were collected on FL1 and FL2 channels, respectively. In each measurement, a minimum of 20000 cells were analysed. Data were acquired and analysed using the Cell Quest software (Becton Dickinson). Relative change in the mean fluorescence intensity was calculated as the ratio between mean fluorescence intensity in the channel of the treated cells and that of the control cells.

### Southern-blot hybridization

For Southern-blot analysis, 5 μg of total DNA of the treated and control cells were linearized by digestion with *Bam*HI (Boehringer Mannheim, Mannheim, Germany), and the digested DNA was electrophoresed at 50 V for 12 h in a 1.0% agarose gel. DNA bands in the gel were denatured and transferred on to a Hybond N<sup>+</sup> nylon membrane (Amersham, Little Chalfont, Bucks., U.K.). Hybridization with a [ $\alpha$ -<sup>32</sup>P]dCTP-labelled human mtDNA probe of the D-loop region (spanning nucleotide positions 16455–1462) was carried out at 65 °C for 2 h in a rapid hybridization buffer (Amersham). The filter was washed with 2 × SSC/0.1% SDS



**Figure 3** Increase of mitochondrial mass in MRC-5 cells treated with H<sub>2</sub>O<sub>2</sub>

(**A**) MRC-5 cells were treated with 180 μM H<sub>2</sub>O<sub>2</sub>. After 48 h of treatment, the mitochondrial mass of the cells stained with NAO was analysed by flow cytometry. The area under the thin (left-hand) curve represents the population of control cells, and the area under the thick (right-hand) curve represents the population of H<sub>2</sub>O<sub>2</sub>-treated cells. (**B**) MRC-5 cells were treated for 24–72 h with 90–360 μM H<sub>2</sub>O<sub>2</sub>, and the relative NAO intensity of treated cells was analysed by flow cytometry. Results are expressed graphically as percentages of the fluorescence intensity of the untreated control (C) cells. Values are means ± S.D. of results from six independent experiments performed in triplicate. Statistical analysis showed that the differences in the NAO intensity between the untreated and H<sub>2</sub>O<sub>2</sub>-treated cells were statistically significant (\*,  $P < 0.05$  versus control).

(where 1 × SSC is 0.15 M NaCl/0.015 M sodium citrate) for 15 min at room temperature and then washed twice with 0.1 × SSC/0.1% SDS for 15 min at 65 °C. The washed filter was wrapped with Saran Wrap ('clingfilm') and subjected to autoradiography.

### MTT reduction assay

Cells were treated with H<sub>2</sub>O<sub>2</sub> at the concentrations from 90 to 360 μM as described above. After 48 h of incubation, cells were harvested and cell viability was determined by the Trypan Blue-exclusion method. For determination of mitochondrial function, we incubated the cells with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT; Sigma) at a final concentration of 2.0 μg/ml during the last 2 h of the incubation period. The cells were lysed with DMSO and the absorbance at 570 nm was measured. An MTT reduction index was calculated


as the ratio between the absorbance at 570 nm and the number of viable cells [27].

### Statistics

Data are presented as means  $\pm$  S.D. except where indicated. Comparisons among multiple groups were made by a one-way analysis of variance (ANOVA) followed by Fisher's protected least significant difference *post hoc* test. The differences with  $P < 0.05$  were considered significant.


### RESULTS

After treatment for 24–72 h with 90–360  $\mu$ M  $H_2O_2$ , growth kinetics and viability of MRC-5 cells were affected in a dose-dependent manner as shown in Figure 1(A) and 1(B), respectively. We found that the untreated cells and the cells treated with 90  $\mu$ M  $H_2O_2$  kept growing steadily, but growth retardation was observed in the cells treated with 180 or 360  $\mu$ M  $H_2O_2$ . Moreover, the rate of increase in the number of the cells treated with 90 or 180  $\mu$ M  $H_2O_2$  was significantly lower than that of the untreated cells ( $P < 0.05$ , Figure 1A). Most of the cells treated with 180 or



**Figure 4** Increase of mitochondrial mass in MRC-5 cells treated with BSO

After treatment of MRC-5 cells with 1 or 2 mM BSO, the production of ROS in the cells, monitored by DCF fluorescence, increased with time (A). After 72 h of treatment, the mitochondrial mass of the cells stained with NAO was analysed by flow cytometry (B). Results are expressed graphically as percentages of the fluorescence intensity of the untreated (C) cells. Values are means  $\pm$  S.D. of the results from two independent experiments performed in triplicate. Statistical analysis showed that the differences in the DCF or NAO intensity between the untreated and BSO-treated cells were statistically significant (\*,  $P < 0.05$  versus control).




**Figure 5** Increase of mtDNA content in MRC-5 cells treated with  $H_2O_2$

After treatment of MRC-5 cells with 180  $\mu$ M  $H_2O_2$  for 24–72 h, the relative mtDNA content of the  $H_2O_2$ -treated cells (H) was determined by Southern-blot hybridization (A). The DNA band intensities were analysed by scanning densitometry and mtDNA content was corrected for variation in sample loading by normalization with the DNA band intensity of the 18 S rRNA gene. (B) Results are expressed graphically as percentages of the mtDNA band intensity of the untreated (C) cells. Values are means  $\pm$  S.D. of the results from three independent experiments. Statistical analysis showed that the differences in the mtDNA band intensity between the untreated and  $H_2O_2$ -treated cells were statistically significant (\*,  $P < 0.05$  versus control).

360  $\mu$ M  $H_2O_2$  appeared enlarged, flattened, and less spindle-shaped as compared with the control cells at the end of 24-h treatment (results not shown). We also found that the viability of the cells treated with 180 or 360  $\mu$ M  $H_2O_2$  was lower than that of the control cells ( $P < 0.05$ ), but still remained 80 % after 72-h treatment (Figure 1B). Furthermore, we found that 24 h after treatment the percentage of cells in  $G_0$  and  $G_1$  phases was increased (Figure 2A), and that the percentage of cells in the S phase decreased as a function of  $H_2O_2$  concentration (Figure 2B).

In an attempt to determine whether  $H_2O_2$  leads to the proliferation of mitochondria, we used the fluorescent dye NAO to monitor the mitochondrial mass [22–24]. After treatment for 24–72 h with the indicated concentrations of  $H_2O_2$ , the relative NAO intensity of the treated cells was found to be significantly higher than that of the untreated cells ( $P < 0.05$ , Figure 3). The relative NAO intensity of the treated cells was increased to 1.3–2.5-fold as compared with that of the control cells, depending on the  $H_2O_2$  concentration and the duration of incubation (Figure 3). We observed a similar increase in the relative intensity of another fluorescent dye, MitoTracker Green FM (results not shown). These results indicate that the oxidative stress induced by  $H_2O_2$  can lead to an increase in the number of mitochondria in MRC-5 cells in a concentration- and time-dependent manner.




**Figure 6** Membrane potential is normal in the extra mitochondria of the  $\text{H}_2\text{O}_2$ -treated cells

After treatment of MRC-5 cells with 90 or 180  $\mu\text{M}$   $\text{H}_2\text{O}_2$  for 48 h, the mitochondrial mass and membrane potential were measured by flow cytometry after staining the cells with NAO and Rhodamine 123 (Rh123), respectively. The relative intensity is expressed graphically as a percentage of the fluorescence intensity of the untreated cells. The increase in mitochondrial membrane potential and mitochondrial mass were correlated throughout the 48-h period of the experiment ( $r = 0.982$ ,  $P < 0.0001$ ).

In addition to the oxidative stress induced by  $\text{H}_2\text{O}_2$ , BSO, an inhibitor of the glutathione biosynthetic pathway, was used to treat the cultured cells. After treatment with 1 or 2 mM BSO, we found that the concentration of ROS in the cells, as revealed by staining with DCFH-DA, increased with time ( $P < 0.05$ , Figure 4A). After treatment for 72 h with 1 or 2 mM BSO, the relative intensity of NAO of the treated cells was increased to 1.6- or 2.2-fold as compared with that of the control cells, respectively ( $P < 0.05$ , Figure 4B). These results indicate that not only  $\text{H}_2\text{O}_2$  but also glutathione depletion can lead to an increase in the number of mitochondria in MRC-5 cells.

Using Southern-blot hybridization, we determined the mtDNA content of the MRC-5 cells after treatment with 180  $\mu\text{M}$   $\text{H}_2\text{O}_2$  for 24–72 h. We found that the mtDNA content of the treated cells was higher than that of the untreated cells and increased with the duration of incubation (Figure 5). This increase of mtDNA content was consistent with the observation of proliferation of mitochondria in the  $\text{H}_2\text{O}_2$ -treated cells as revealed by NAO staining.

To analyse whether the extra mitochondria in the  $\text{H}_2\text{O}_2$ -treated cells were functional, we measured mitochondrial membrane potential of the treated cells by staining with Rhodamine 123 [25]. The relative intensity of Rhodamine 123 of the  $\text{H}_2\text{O}_2$ -treated cells was significantly higher than that of the control cells and was increased with the duration of incubation (results not shown). The increase in mitochondrial membrane potential correlated well with the increase in mitochondrial mass throughout the 48 h period of the experiment ( $r = 0.982$ ,  $P < 0.0001$ , Figure 6). The mitochondrial membrane potential per unit mitochondrial mass, expressed as the ratio of fluorescence intensity of Rhodamine 123 to that of NAO, appeared to be similar throughout the 48 h of the experiment (Figure 6). Moreover, we observed similar results using another fluorescent dye, DiOC<sub>6</sub>(3), the uptake of which is sensitive to mitochondrial membrane potential (results not shown).




**Figure 7** The increased mitochondria in the  $\text{H}_2\text{O}_2$ -treated cells are functional

Cells were treated with  $\text{H}_2\text{O}_2$  at concentrations from 90 to 360  $\mu\text{M}$  as described in the text. After 48 h of incubation, the capacity of the treated cells to reduce MTT was analysed as described in the Materials and methods section. The absorbance at 570 nm of the formosan formed in the MTT assay (●) and the number of viable cells (○) were determined concurrently after incubation of the MRC-5 cells for 48 h with the indicated concentrations of  $\text{H}_2\text{O}_2$ , and are expressed as percentages of the control (A). MTT reduction index was calculated as the ratio between the absorbance at 570 nm and the number of viable cells counted using a haemocytometer after 48 h of incubation with indicated concentrations of  $\text{H}_2\text{O}_2$  (B). The data are expressed as the percentage of controls (C; means  $\pm$  S.D.) from two independent experiments in triplicate. Statistical analysis showed that the differences in MTT reduction index between  $\text{H}_2\text{O}_2$ -treated and control cells are statistically significant (\*,  $P < 0.05$  versus control).


For the assessment of mitochondrial function, we analysed the capacity of  $\text{H}_2\text{O}_2$ -treated cells to reduce MTT. We found that although the number of the viable cells after  $\text{H}_2\text{O}_2$  treatment was significantly decreased ( $P < 0.05$ ), the absorbance at 570 nm of the remaining cells was only slightly decreased as compared with the untreated cells (Figure 7A). An arbitrary MTT reduction index, calculated as the ratio between the absorbance at 570 nm and the corresponding number of viable cells, revealed a significant increase in the relative MTT activity ( $P < 0.05$ , Figure 7B). These results indicate that the mitochondria increased in the  $\text{H}_2\text{O}_2$ -treated cells were functional.

To test whether the alteration of mitochondrial mass and mtDNA copy number in the cell might be uncoupled from the cell-cycle-controlled biosynthesis of chromosomal DNA, we treated MRC-5 cells with mimosine, lovastatin or genistein,



**Figure 8** Cell-cycle arrest induced by mimosine, lovastatin and genistein

MRC-5 cells were treated with 200  $\mu$ M mimosine, 2.5  $\mu$ M lovastatin and 50  $\mu$ M genistein in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum. After 24 h of incubation at 37 °C, the cell cycle was analysed on a flow cytometer as described in the Materials and methods section. Typical results from untreated cells (A) and cells treated with mimosine (B), lovastatin (C) and genistein (D) are presented.



**Figure 9** Increase of mitochondrial mass in MRC-5 cells treated with drugs affecting the cell cycle

MRC-5 cells were treated with 200  $\mu$ M mimosine, 2.5  $\mu$ M lovastatin and 50  $\mu$ M genistein to cause cell-cycle arrest. After 48 h of incubation at 37 °C, the relative NAO intensity of treated cells was analysed by flow cytometry as described in the Materials and methods section. The results are expressed graphically as a percentage of the fluorescence intensity of the untreated MRC-5 cells. Values are means  $\pm$  S.D. of the results obtained from three independent experiments. Statistical analysis showed that the differences in the NAO intensity between the untreated and drug-treated cells were statistically significant (\*,  $P < 0.05$  versus control).

which cause cell-cycle arrest. We found that after 24 h of treatment the cell cycle was arrested at the late G<sub>1</sub>, G<sub>0</sub>/G<sub>1</sub> and G<sub>2</sub>/M phases by mimosine, lovastatin and genistein, respectively

(Figure 8). Moreover, after 48 h of treatment the relative intensity of NAO of the treated cells was increased to 2.3-, 1.7- and 2.1-fold, respectively, as compared with that of the control cells ( $P < 0.05$ , Figure 9).

## DISCUSSION

Each human and animal cell contains between several hundred and over a thousand mitochondria, each carrying 2–10 copies of mtDNA. The mitochondrial mass and mtDNA copy number of individual cells vary with the type of the cell and tissue, and are changed during cell differentiation, hormone treatment and exercise [28–31].

In the present study, we observed growth arrest in human lung fibroblasts treated with low concentrations of H<sub>2</sub>O<sub>2</sub> (Figures 1 and 2). It is apparent that H<sub>2</sub>O<sub>2</sub> at concentrations of between 90 and 180  $\mu$ M increased the population-doubling time and that 360  $\mu$ M H<sub>2</sub>O<sub>2</sub> caused a permanent growth arrest. Most of the treated cells exhibited significant alterations in morphology as compared with the control cells. The results are consistent with the previous observations that H<sub>2</sub>O<sub>2</sub> at sublethal concentrations can induce senescence-like growth arrest in human cells [32,33]. Moreover, we found in this study that the oxidative stress induced by sublethal concentrations of H<sub>2</sub>O<sub>2</sub> caused an increase in the amount of mitochondria and mtDNA of the treated cells and that the increased mitochondria were functional. Furthermore, the oxidative stress induced by reduction of the intracellular glutathione level with BSO was also found to increase mitochondria in the treated cells (Figure 4). In a separate experiment,

we exposed MRC-5 cells to 20  $\mu$ M antimycin A and observed that the ROS generation in the treated cells was increased 2-fold compared with that of the untreated cells after 30 min of treatment. Moreover, after treatment for 72 h, we found that the relative intensity of NAO in the treated cells was increased to 2.6-fold compared with that of the control cells. This is consistent with the observation that by partial inhibition of respiratory function by antimycin A, the expressions of both cytochrome *c*<sub>1</sub> and cytochrome *b* are increased in human fibroblasts [34]. Moreover, intracellular H<sub>2</sub>O<sub>2</sub> level was found to increase within 3 h prior to expression of these genes. These results indicate together that mild oxidative stress can trigger cell-cycle arrest and lead to an increase of mitochondrial mass and mtDNA in human cells.

Increases of mitochondria and mtDNA synthesis have been observed to occur in the absence of nuclear DNA replication [35]. The alteration of mitochondrial mass and mtDNA in the cell might be uncoupled from the cell-cycle-controlled biosynthesis of chromosomal DNA. To confirm this, we treated MRC-5 cells with lovastatin, genistein or mimosine and observed that these drugs caused cell-cycle arrest (Figure 8). In addition, the mitochondrial mass was increased, to a different extent, in the cells after 48 h of treatment (Figure 9). It is worth noting that the amplitude of increase of the mitochondrial mass in the cells treated with the cell-cycle-arresting drugs was similar to that in the H<sub>2</sub>O<sub>2</sub>-treated cells within the same treatment period. The results imply that whereas overall cell division was arrested by H<sub>2</sub>O<sub>2</sub> or the drugs, mitochondrial biogenesis was unaffected and that mitochondria continued to proliferate as if the cells were going to divide. Thus, mild oxidative stress may induce an increase of mitochondria and mtDNA via a pathway that bypasses cell-cycle control.

Increase of mitochondria has also been observed in some tumour cells treated with herbimycin A and genistein [24,27]. Mancini et al. [24] found that herbimycin A, a tyrosine kinase inhibitor, induces cell-cycle arrest in G<sub>2</sub>/M and results in a marked accumulation of mitochondria in a human colon carcinoma cell line (Colo-205). In addition, the G<sub>2</sub>/M cell-cycle arrest and increase of mitochondrial number and/or function were also observed in three tumour cell lines (MCF-7, human breast tumour; Jurkat cells, human T-cell leukaemia; and L-929, mouse transformed fibroblasts) after treatment of genistein [27]. The results obtained in these studies are consistent with our findings and support the notion that the cell-cycle arrest induced by oxidative stress plays a critical role in the increase of mitochondria and mtDNA.

Furthermore, it has been found that the relative content of mtDNA is increased with age and that the mtDNA content is correlated with the level of oxidative damage to DNA in human lung tissues [4]. Both oxidative DNA damage and lipid peroxidation are enhanced during aging in the human lung [4–7], indicating that there is a higher oxidative stress in the tissues of aged individuals. Therefore, the endogenous and/or exogenous oxidative stress in the lung of elderly subjects may cause cell-cycle arrest and thereby increase the amount of mitochondria and mtDNA in lung tissues.

In summary, we have demonstrated clearly for the first time that oxidative stress induced by H<sub>2</sub>O<sub>2</sub> leads to an increase of mitochondria and mtDNA in human lung fibroblasts. Moreover, the mitochondria increased by H<sub>2</sub>O<sub>2</sub>-elicited oxidative stress are functional. We therefore suggest that the increase in mitochondrial mass and mtDNA content may be one of the early molecular events gearing the human cells up to respond to endogenous or exogenous oxidative stress through cell-cycle arrest.

This work was supported by a grant (NSC89-2316-B010-012) from the National Science Council and a 5-year national laboratory grant (NHRI-GT-EX89S05L) from the National Health Research Institutes, Republic of China.

## REFERENCES

- Chance, B., Sies, H. and Boveris, A. (1979) Hydroperoxide metabolism in mammalian organs. *Physiol. Rev.* **59**, 527–604
- Papa, S. (1996) Mitochondrial oxidative phosphorylation changes in the life span. Molecular aspects and physiopathological implications. *Biochim. Biophys. Acta* **1276**, 87–105
- Wei, Y. H. (1998) Oxidative stress and mitochondrial DNA mutations in human aging. *Proc. Soc. Exp. Biol. Med.* **217**, 53–63
- Lee, H. C., Lu, C. Y., Fahn, H. J. and Wei, Y. H. (1998) Aging- and smoking-associated alteration in the relative content of mitochondrial DNA in human lung. *FEBS Lett.* **441**, 292–296
- Fahn, H. J., Wang, L. S., Hsieh, R. H., Chang, S. C., Kao, S. H., Huang, M. H. and Wei, Y. H. (1996) Age-related 4,977 bp deletion in human lung mitochondrial DNA. *Am. J. Respir. Crit. Care Med.* **154**, 1141–1145
- Fahn, H. J., Wang, L. S., Kao, S. H., Chang, S. C., Huang, M. H. and Wei, Y. H. (1998) Smoking-associated mitochondrial DNA mutations and lipid peroxidation in human lung tissues. *Am. J. Respir. Cell Mol. Biol.* **19**, 901–909
- Lee, H. C., Lim, M. L. R., Lu, C. Y., Liu, V. W. S., Fahn, H. J., Zhang, C., Nagley, P. and Wei, Y. H. (1999) Concurrent increase of oxidative DNA damage and lipid peroxidation together with mitochondrial DNA mutations in human lung tissues during aging-smoking enhances oxidative stress on the aged tissues. *Arch. Biochem. Biophys.* **362**, 309–316
- Wei, Y. H., Kao, S. H. and Lee, H. C. (1996) Simultaneous increase of mitochondrial DNA deletions and lipid peroxidation in human aging. *Ann. N.Y. Acad. Sci.* **786**, 24–43
- Lu, C. Y., Lee, H. C., Fahn, H. J. and Wei, Y. H. (1999) Oxidative damage elicited by imbalance of free radical scavenging enzymes is associated with large-scale mtDNA deletions in aging human skin. *Mutat. Res.* **423**, 11–21
- Gadaleta, M. N., Rainaldi, G., Lezza, A. M. S., Milella, F., Fracasso, F. and Cantatore, P. (1992) Mitochondrial DNA copy number and mitochondrial DNA deletion in adult and senescent rats. *Mutat. Res.* **275**, 181–193
- Barrientos, A., Casademont, J., Cardellach, F., Estivill, X., Urbano-Marquez, A. and Nunes, V. (1997) Reduced steady-state levels of mitochondrial RNA and increased mitochondrial DNA amount in human brain with aging. *Mol. Brain Res.* **52**, 284–289
- Ames, B. N., Shigenaga, M. K. and Hagen, T. M. (1995) Mitochondrial decay in aging. *Biochim. Biophys. Acta* **1271**, 165–170
- Sohal, R. S. and Sohal, B. H. (1991) Hydrogen peroxide release by mitochondria increases during aging. *Mech. Ageing Dev.* **57**, 187–202
- Richter, C., Gogvadze, V., Laffranchi, R., Schlapbach, R., Schnizer, M., Suter, M., Walter, P. and Yaffee, M. (1995) Oxidants in mitochondria: from physiology to disease. *Biochim. Biophys. Acta* **1271**, 67–74
- Sohal, R. S., Sohal, B. H. and Orr, W. C. (1995) Mitochondrial superoxide and hydrogen peroxide generation, protein oxidative damage and longevity in different species of flies. *Free Radical Biol. Med.* **19**, 499–504
- Sen, C. K. and Packer, L. (1996) Antioxidant and redox regulation of gene transcription. *FASEB J.* **10**, 709–720
- Lander, H. M. (1997) An essential role for free radicals and derived species in signal transduction. *FASEB J.* **11**, 118–124
- Burdon, R. H. (1995) Superoxide and hydrogen peroxide in relation to mammalian cell proliferation. *Free Radical Biol. Med.* **18**, 775–794
- Poyton, R. O. and McEwen, J. E. (1996) Crosstalk between nuclear and mitochondrial genomes. *Annu. Rev. Biochem.* **65**, 563–607
- Scarpulla, R. C. (1997) Nuclear control of respiratory chain expression in mammalian cells. *J. Bioenerg. Biomembr.* **29**, 109–119
- Lee, H. C. and Wei, Y. H. (2000) Mitochondrial role in life and death of the cell. *J. Biomed. Sci.* **7**, 2–15
- Maftah, A., Petit, J. M., Ratinaud, M. H. and Julien, R. (1989) 10-N nonyl acridine orange: a fluorescent probe which stains mitochondria independently of their energetic state. *Biochem. Biophys. Res. Commun.* **164**, 185–190
- Petit, J. M., Maftah, A., Ratinaud, M. H. and Julien, R. (1992) 10-N nonyl aridine orange interacts with cardiolipin and allows the quantification of this phospholipid in isolated mitochondria. *Eur. J. Biochem.* **209**, 267–273
- Mancini, M., Anderson, B. O., Caldwell, E., Sedghinasab, M., Paty, P. B. and Hockenbery, D. M. (1997) Mitochondrial proliferation and paradoxical membrane depolarization during terminal differentiation and apoptosis in a human colon carcinoma cell line. *J. Cell Biol.* **138**, 449–469
- Rothe, G. and Valet, G. (1990) Flow cytometric analysis of respiratory burst activity in phagocytes with hydroethidine and 2',7'-dichlorofluorescein. *J. Leuk. Biol.* **47**, 440–448

- 26 Garcia-Ruiz, C., Colell, A., Morales, A., Kaplowitz, N. and Fernandez-Checa, J. C. (1995) Role of oxidative stress generated from mitochondrial electron transport chain and mitochondrial glutathione status in loss of mitochondrial function and activation of transcription factor nuclear factor- $\kappa$ B: studies with isolated mitochondria and rat hepatocytes. *Mol. Pharmacol.* **48**, 825–834
- 27 Pagliacci, M. C., Spinozzi, F., Migliorati, G., Fumi, G., Smacchia, M., Grignani, F., Riccardi, C. and Nicoletti, I. (1993) Genistein inhibits tumour cell growth *in vitro* but enhances mitochondrial reduction of tetrazolium salts: a further pitfall in the use of the MTT assay for evaluating cell growth and survival. *Eur. J. Cancer* **29A**, 1573–1577
- 28 Robin, E. D. and Wong, R. (1988) Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells. *J. Cell. Physiol.* **136**, 507–513
- 29 Renis, M., Cantatore, P., Polosa, P. L., Fracasso, F. and Gadaleta, M. N. (1989) Content of mitochondrial DNA and of three mitochondrial RNAs in developing and adult cerebellum. *J. Neurochem.* **52**, 750–754
- 30 Shay, J. W., Pierce, D. J. and Werbin, H. (1990) Mitochondrial DNA copy number is proportional to total cell DNA under a variety of growth conditions. *J. Biol. Chem.* **265**, 14802–14807
- 31 Williams, R. S., Salmons, S., Newsholme, E. A., Kaufman, R. E. and Mellor, J. (1986) Regulation of nuclear and mitochondrial gene expression by contractile activity in skeletal muscle. *J. Biol. Chem.* **261**, 376–380
- 32 Bladier, C., Wolvetang, E. J., Hutchinson, P., de Haan, J. B. and Kola, I. (1997) Response of a primary human fibroblast cell line to H<sub>2</sub>O<sub>2</sub>: senescence-like growth arrest or apoptosis. *Cell Growth Differ.* **8**, 589–598
- 33 Chen, Q. M., Bartholomew, J. C., Campisi, J., Acosta, M., Reagan, J. D. and Ames, B. N. (1998) Molecular analysis of H<sub>2</sub>O<sub>2</sub>-induced senescent-like growth arrest in normal human fibroblasts: p53 and Rb control G<sub>1</sub> arrest but not cell replication. *Biochem. J.* **332**, 43–50
- 34 Suzuki, H., Kumagai, T., Goto, A. and Sugiura, T. (1998) Increase in intracellular hydrogen peroxide and upregulation of a nuclear respiratory gene evoked by impairment of mitochondrial electron transfer in human cells. *Biochem. Biophys. Res. Commun.* **249**, 542–545
- 35 Sazer, S. and Sherwood, S. W. (1990) Mitochondrial growth and DNA synthesis occur in the absence of nuclear DNA replication in fission yeast. *J. Cell Sci.* **97**, 509–516

Received 30 September 1999/8 March 2000; accepted 29 March 2000