



Power training in elite young soccer players: Effects of using loads above or below the optimum power zone

Irineu Loturco, Lucas A. Pereira, Valter P. Reis, Chris Bishop, Vinicius Zanetti,
Pedro E. Alcaraz, Tomás T. Freitas & Michael R. McGuigan

To cite this article: Irineu Loturco, Lucas A. Pereira, Valter P. Reis, Chris Bishop, Vinicius Zanetti, Pedro E. Alcaraz, Tomás T. Freitas & Michael R. McGuigan (2019): Power training in elite young soccer players: Effects of using loads above or below the optimum power zone, Journal of Sports Sciences, DOI: [10.1080/02640414.2019.1651614](https://doi.org/10.1080/02640414.2019.1651614)

To link to this article: <https://doi.org/10.1080/02640414.2019.1651614>



Published online: 07 Aug 2019.



Submit your article to this journal [↗](#)






View Crossmark data [↗](#)

ORIGINAL ARTICLE



Power training in elite young soccer players: Effects of using loads above or below the optimum power zone

Irineu Loturco ^{a,b,i}, Lucas A. Pereira^a, Valter P. Reis^a, Chris Bishop^c, Vinicius Zanetti^d, Pedro E. Alcaraz ^{e,f}, Tomás T. Freitas ^e and Michael R. McGuigan^{g,h}

^aNAR - Nucleus of High Performance in Sport, São Paulo, Brazil; ^bDepartment of Human Movement Science, Federal University of São Paulo, São Paulo, Brazil; ^cFaculty of Science and Technology, London Sports Institute, Middlesex University, London, UK; ^dRed Bull Brazil Football, Jarinú, Brazil; ^eResearch Center for High Performance Sport, Catholic University of Murcia, Murcia, Spain; ^fFaculty of Sport Sciences, Catholic University of Murcia, Murcia, Spain; ^gSports Performance Research Institute New Zealand (SPRINZ), Auckland University of Technology, Auckland, New Zealand; ^hSchool of Medical and Health Sciences, Edith Cowan University, Perth, Australia; ⁱUniversity of South Wales, Pontypridd, Wales, UK

ABSTRACT

This study aimed to examine the effects of two jump squat (JS) training programs involving different loading ranges in under-20 soccer players during a preseason period. Twenty-three elite young soccer players performed sprint speed (at 5-, 10-, and 20-m), change-of-direction (COD) speed, JS peak-power (PP), and countermovement jump (CMJ) tests pre and post four weeks of training. Athletes were pair-matched in two groups according to their optimum power loads (OPL) as follows: lower than OPL (LOPL; athletes who trained at a load 20% lower than the OPL) and higher than OPL (HOPL; athletes who trained at a load 20% higher than the OPL). Magnitude-based inferences were used to compare pre- and post-training measures. Meaningful increases in the PP JS were observed for both groups. Likely and possible improvements were observed in the 5- and 10-m sprint velocity in the LOPL group. Meanwhile, possible and likely improvements were observed in the CMJ, 5- and 10-m sprint velocity, and COD speed in the HOPL group. Overall, both training schemes induced positive changes in athletic performance. Soccer coaches and sport scientists can implement the JS OPL-based training schemes presented here, either separately or combined, to improve the physical performance of youth soccer players.

ARTICLE HISTORY

Accepted 29 July 2019

KEYWORDS

Team-sports; football; speed ability; vertical jump; optimal loads

Introduction

Improving speed and power performance during professional soccer preseasons has long been considered a major challenge for coaches and sport scientists (Loturco, Pereira, & Kobal et al., 2015; Meckel, Harel, Michaely, & Eliakim, 2014; Mercer, Gleeson, & Mitchell, 2014). This issue is typically associated with the well-established concurrent training effects, which appear to hamper the adequate development of neuromuscular capacities in periods where high volumes of aerobic exercise (e.g., technical and tactical workouts) are applied (Docherty & Sporer, 2000; Helgerud, Rodas, Kemi, & Hoff, 2011; Kraemer, French, & Paxton et al., 2004; Loturco et al., 2015). For some authors, the interference between endurance, speed, and power adaptations can be explained by several factors such as: 1) the inability of muscle to adapt to distinct stimuli due to simultaneous requirements from different metabolic pathways; 2) residual fatigue induced by successive training sessions; 3) age, individual training background, and physiological traits; and 4) the type of resistance training program (Noon, James, Clarke, Akubat, & Thake, 2015; Tufano, Brown, & Haff, 2017). Among these aspects, the latter is the only one that practitioners can manipulate in certain ways.

More recently, the optimum power load (OPL) has been used as a practical and effective alternative to improve speed and power performance in elite soccer players (Loturco, Nakamura, & Kobal et al., 2016; Loturco, Pereira, & Kobal

et al., 2016). The “optimum power zone” can be defined as the range of loads able to maximize power output in some resistance exercises (Loturco, Nakamura, & Tricoli et al., 2015). This mechanical phenomenon usually occurs at light or moderate loading conditions (i.e., ~30–70% one-repetition maximum [1RM]), and varies according to the lift in question (e.g., bench press or half squat) and its respective mode of execution (e.g., traditional or ballistic) (Cronin, McNair, & Marshall, 2001; Kawamori & Haff, 2004; Loturco, Pereira, Kobal, & McGuigan, 2019). The OPL is typically found at a narrow range of bar-velocities, independent of subjects’ training background, sport discipline, and strength-power level (Loturco, Kobal, & Moraes et al., 2017; Loturco et al., 2015; Sanchez-Medina, Perez, & Gonzalez-Badillo, 2010). Importantly, it has been reported that this load is capable of improving the physical capacities at both ends of the force-velocity curve (i.e., high force, low velocity portion; low force, high velocity portion) and counteracting the speed-power decrements which normally occur in response to congested soccer pre-seasons (Loturco, Kobal, & Kitamura et al., 2017; Loturco et al., 2015; Meckel et al., 2014; Mercer et al., 2014; Taylor, Portas, Wright, & Weston, 2012). However, it is still unknown how the power-load relationship is affected when athletes train immediately below or above the optimum training intensity (e.g., using loads 20% higher or lower than the OPL).

In this context, it has been suggested that training with lower loads and higher velocities might lead to greater adaptations in speed qualities, whereas training with higher loads and lower velocities would result in superior gains in strength-related performance (Behm & Sale, 1993; Coyle, Feiring, & Rotkis et al., 1981; Cronin et al., 2001; Kanehisa & Miyashita, 1983; Cormie, McGuigan, & Newton, 2011b). Accordingly, in a study with soccer players who trained under different loading conditions for 6 weeks (i.e., “reduced velocity group” [RVG] and “increased velocity group” [IVG]), the authors detected higher increases in leg press 1RM in the RVG. In contrast, greater improvements in linear and change of direction (COD) speed were noted for the IVG (Loturco, Nakamura, & Kobal et al., 2015). Similarly, McBride, Triplett-McBride, Davie, and Newton (2002) compared the effects of an 8-week training program with heavy- (80% 1RM) versus light-load (30% 1RM) jump squats (JS) on various physical measures, observing an overall trend toward enhanced velocity capabilities (e.g., 10-m sprint time, peak power [PP], and peak velocity at 30% 1RM) in the light-load group. On the other hand, the heavy-load group showed significant improvements in PP and peak force (only) at heavier loading conditions (i.e., 55–80% 1RM) and, remarkably, presented a significant and unexpected decrease in sprint performance over very-short distances (i.e., 5-m) (which also supports the concept of velocity-specificity in strength-power training) (Cormie et al., 2011b).

Therefore, it is important to establish an upper (and also a lower) limit of loads capable of eliciting positive changes in both speed and power-related capabilities. This is particularly relevant in elite soccer, where straight sprinting and explosive actions (e.g., vertical jumps) play a crucial role, being directly related to decisive game situations (i.e., scoring or assisting a goal) (Faude, Koch, & Meyer, 2012). Considering the aforementioned challenges and the effectiveness of OPL in promoting positive adaptations and reducing the possible impairments in speed-power performance during high-volume soccer preseasons (Loturco et al., 2015), it is reasonable to use this range of loads as a basis for defining the inferior and superior power-training zones. The aim of this study was to examine the effects of two different JS training programs (using loads 20% higher or 20% lower than the OPL) on the athletic performance (e.g., linear speed, COD speed, and loaded and unloaded jumping ability) of elite young soccer players during a preseason period.

Methods

Participants

Twenty-three male under-20 players from the same soccer club with at least six years of experience in a professional academy (age: 18.3 ± 0.7 years, ranging between 18 and 19 years; height: 178.3 ± 5.4 cm; body-mass [BM]: 71.5 ± 6.5 kg) regularly competing in the most important regional Brazilian youth tournament took part in this study. Athletes were pair-matched in two training groups according to the load associated with maximum PP output (i.e., OPL) in the JS exercise as follows: lower than optimum power load (LOPL, $n = 12$; athletes who trained at a load 20% lower than the OPL) and higher than optimum power load (HOPL, $n = 11$; athletes who trained at a load 20% higher than the OPL). The study protocol took place during a four-week pre-season training phase, after a four-week period without any programmed training sessions. The study was approved by the local Ethics Committee and the participants signed an informed consent form prior to research commencement.

Study design

A parallel two-group, randomized, longitudinal design was conducted to test the effectiveness of two distinct training programs on the neuromuscular performance of elite young soccer players during a four-week pre-season training period (Figure 1). Players were grouped in pairs according to the baseline results of their PP output in the JS, and subsequently the group allocation was performed by tossing a coin. All athletes had been previously familiarized with the performance tests, which were performed in the following order: countermovement jump (CMJ), sprinting speed at 5-, 10-, and 20-m, COD speed, and PP JS. The physical tests were performed on the same day, both pre- and post-training. Prior to all testing sessions, a general and specific warm-up routine was performed, involving light running (5-min at a self-selected pace) and submaximal attempts at each testing exercise (e.g., submaximal sprints and vertical jumps).

Training program

During the experimental period, all soccer players performed 12 power-oriented training sessions. The players involved in this study participated in all power training sessions during the pre-season training period. A typical weekly training




Figure 1. Schematic presentation of the study design. CMJ: countermovement jump; VEL: sprint velocity; PP: peak power; JS: jump squat exercise; OPL: optimum power load; LOPL: lower than OPL group; HOPL: higher than OPL group.

Table 1. Typical weekly training program for the soccer players during the 4 weeks of preseason.

	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning	PT 30'	TEC/TAC 60'	PT 30'	Rest	PT 30'	TEC/TAC 70'
Afternoon	TEC/TAC 70'	TEC/TAC 80'	TEC/TAC 90'	TEC/TAC 80'	TEC/TAC 90'	Rest

Note: TEC = Technical Training; TAC = Tactical Training; PT = Power Training; The numbers after the training sessions represent the volume in minutes. TEC/TAC training involved different formats of small-sided games and specific technical actions (e.g., goal shooting, corner kick situations).

schedule is presented in Table 1. The power training sessions consisted of performing 6 sets of 6 repetitions of the JS exercise at a load corresponding to either 20% lower than the OPL (LOPL group) or 20% higher than the OPL (HOPL group). These loading intensities were chosen because at $\pm 20\%$ of the OPL, athletes usually produce $\sim 90\%$ of their maximum power output in the JS exercise, which can still be considered a substantial amount of power. For both groups, the training loads were controlled and adjusted every four training sessions according to the OPL-based values, as follows: (sessions 1–4) OPL; (sessions 5–8) 1.05 x OPL; (sessions 9–12) 1.10 x OPL (Loturco et al., 2015).

Testing procedures

Vertical jumping test

Vertical jump height was determined using the CMJ. The soccer players were instructed to execute a downward movement followed by complete extension of the legs. All attempts were executed with the hands placed on the hips. The CMJ was performed on a contact platform (Elite Jump System®; S2 Sports, São Paulo, Brazil). A total of five attempts were allowed, interspersed by 15-s. The best attempt was retained for data analysis purposes.

Peak power in the jump squat exercise

Maximum PP output in the JS was assessed on a Smith machine (Hammer Strength, Rosemont, IL, USA). Players were instructed to execute two repetitions at maximal velocity for each load, starting at 40% of their BM. Athletes executed knee flexion until the thigh was parallel to the ground ($\sim 100^\circ$ knee angle) and, after a command, jumped as fast as possible without losing contact between their shoulder and the bar. A load of 10% BM was gradually added until a decrease in PP was observed. A 5-minute interval between sets was provided. To determine PP, a linear transducer (T-Force, Dynamic Measurement System; Ergotech Consulting S.L., Murcia, Spain) was attached to the Smith machine bar. The load corresponding to the maximum PP value was considered as the OPL and was used as a reference to calculate the loads for both groups of training. The maximum PP values for the loads corresponding to the OPL, 20% lower than the OPL (-20% OPL), and 20% higher than the OPL ($+20\%$ OPL) relative to the players' BM were retained for analysis.

Sprinting speed

Four pairs of photocells (Smart Speed, Fusion Sport, Brisbane, AUS) were positioned at the starting line and at the distances of 5-, 10-, and 20-m. The soccer players sprinted twice, starting from a standing position 0.3-m behind the starting line. The sprint tests were performed on an indoor running track. Sprint velocity (VEL) was calculated as the distance traveled over

a measured time interval. A 5-min rest interval was allowed between the two attempts and the fastest time was considered for subsequent analyses.

Zigzag change of direction speed test

The COD course consisted of four 5-m sections marked with cones set at 100° angles, on an indoor court (Figure 2). Athletes were required to decelerate and accelerate as fast as possible without losing body stability. Two maximal attempts were performed with a 5-min rest interval between attempts. Starting from a standing position with the front foot placed 0.3-m behind the first pair of photocells (i.e., starting line), athletes ran and changed direction as quickly as possible, until crossing the second pair of photocells, placed 20-m from the starting line. The fastest time from the two attempts was retained for analyses.

Statistical analysis

Data are presented as mean \pm standard deviation (SD). To analyze the differences in the CMJ, VEL in all distances tested, COD velocity, and PP JS in both LOPL and HOPL groups, pre- and post-training, the magnitude-based inferences were calculated (Batterham and Hopkins, 2006). The magnitude of the within-group changes in the different performance variables, or between-group differences in the changes, were expressed as standardized mean differences. The smallest worthwhile change was set by using a small effect size ($ES = 0.2$) for each variable tested (Hopkins, Marshall, Batterham, & Hanin, 2009). The quantitative chances of finding differences in the variables tested were assessed qualitatively as follows: $<1\%$, almost certainly not; 1% to 5%, very unlikely; 5% to 25%, unlikely; 25% to 75%, possible; 75% to 95%, likely; 95% to




Figure 2. Schematic presentation of the Zigzag change of direction speed test. The circles represent the positions of the photocells.

99%, very likely; >99%, almost certain. A meaningful difference was considered using the clinical inference, based on threshold chances of harm and benefit of 0.5% and 25% (Hopkins et al., 2009). Additionally, the magnitudes of the standardized differences were interpreted using the following thresholds: <0.2, 0.2–0.6, 0.6–1.2, 1.2–2.0, 2.0–4.0, and >4.0 for trivial, small, moderate, large, very large, and near perfect, respectively (Hopkins et al., 2009). All performance tests used herein demonstrated small errors of measurement, as evidenced by their high levels of accuracy and reproducibility (coefficient of variation <5% and intraclass correlation coefficient >0.90 for all assessments) (Hopkins et al., 2009).

Results

Figure 3 shows the comparisons of the PP outputs in the JS exercise for the different loads tested pre and post the pre-season training period in both training groups. Likely to very likely increases were observed in the PP comparing pre- and post-training measurements in the LOPL group in the three loads analyzed (ES = 0.64, 0.68, and 0.54, for –20% OPL, OPL, and +20% OPL, respectively). Meanwhile, a possible increase was noted in the PP JS in the HOPL group for the OPL and the +20% OPL (ES = 0.23 and 0.48, respectively).




Figure 3. Comparisons of the relative peak power (PP) in the jump squat exercise pre and post the pre-season training period in both training groups. The loads corresponding to the optimum power load (OPL), 20% lower than the OPL (–20% OPL), and 20% higher than the OPL (+20% OPL) were analyzed. LOPL: lower than OPL group; HOPL: higher than OPL group; *possible, #likely, and *very likely within-group effect of time.

Table 2 shows the comparisons of the CMJ height, and sprint and Zigzag velocities pre and post the pre-season training period. A likely and a possible increase in the VEL 5-m and VEL 10-m were detected in the LOPL group, respectively. In the HOPL group, a possible improvement in CMJ height, VEL 5-m, and VEL 10-m was observed, while a likely increase was detected in the COD velocity.

Figure 4 shows the standardized mean differences (ES) for the comparisons of the between-group delta changes in the physical tests performed. No meaningful differences were observed for the CMJ, VEL 5-, 10-, and 20-m, and Zigzag (ES [% chance] = 0.15 [36/63/01], 0.09 [29/30/41], 0.05 [27/38/35], 0.13 [40/47/13], and 0.42 [70/23/7], respectively). In addition, the LOPL group demonstrated higher increases in the PP JS for the –20% OPL and OPL (ES [% chance] = 0.51 [02/15/83] and 0.59 [01/11/88], respectively) in relation to the HOPL, while no meaningful differences were noted in the PP JS for the +20% OPL (ES [% chance] = 0.14 [26/29/45]).

Discussion

The study compared the effects of two different JS training programs (using loads 20% higher or 20% lower than the OPL) in elite young soccer players during a pre-season period. The main findings were: 1) despite the use of lower loads, the LOPL increased power production over the entire range of loads (–20% OPL, OPL, and +20% OPL); 2) the HOPL improved power output only at higher loading conditions (OPL, and +20% OPL); and 3) overall, both training schemes were able to induce positive changes in athletic performance, with meaningful and relevant differences between them.

Despite some controversy regarding this issue, several studies have demonstrated that neuromechanical adaptations are velocity-specific (Behm & Sale, 1993; Coyle et al., 1981; Cronin et al., 2001; Kanehisa & Miyashita, 1983; Cormie et al., 2011b). For example, Brown and Whitehurst (2003) compared the effects of “fast” (4.18 rad·s^{–1}) and “slow” (1.04 rad·s^{–1}) isokinetic training on force and “rate of velocity development”, showing that significant improvements in acceleration occur exclusively at the trained velocity, which, according to the authors, might serve to counterbalance force deficits in power production (when considering the force-velocity relationship). Similarly, a study of under-20 soccer players indicated that increasing bar-velocity during JS (using a system

Table 2. Comparisons of the physical tests pre and post the pre-season training period in under 20 elite soccer players.

	LOPL					HOPL				
	Pre	Post	Δ%	ES (90% CL) rating	% Chance rating	Pre	Post	Δ%	ES (90% CL) rating	% Chance rating
CMJ (cm)	44.9 ± 6.4	45.3 ± 6.4	0.9	0.06 (–0.07; 0.19)	00/96/04 trivial	43.9 ± 4.3	45.1 ± 3.9	2.7	0.26 (0.04; 0.48)	00/32/68 possibly
VEL 5-m (m·s ^{–1})	5.12 ± 0.17	5.24 ± 0.23	2.3	0.64 (–0.20; 1.48)	05/13/82 moderate	5.03 ± 0.34	5.13 ± 0.22	2.0	0.26 (–0.20; 0.73)	05/35/60 possibly
VEL 10-m (m·s ^{–1})	5.91 ± 0.18	5.98 ± 0.26	1.2	0.41 (–0.25; 1.06)	05/23/72 small	5.86 ± 0.27	5.92 ± 0.23	1.0	0.23 (–0.19; 0.65)	04/40/56 possibly
VEL 20-m (m·s ^{–1})	6.84 ± 0.21	6.84 ± 0.26	0	0.03 (–0.34; 0.41)	14/64/22 trivial	6.79 ± 0.25	6.83 ± 0.26	0.6	0.15 (–0.20; 0.51)	06/54/40 unclear
Zigzag (m·s ^{–1})	3.44 ± 0.09	3.45 ± 0.11	0.3	0.04 (–0.62; 0.69)	26/41/33 trivial	3.41 ± 0.11	3.46 ± 0.15	1.5	0.48 (0.03; 0.92)	01/13/86 likely

Note: LOPL: lower than optimum power load group; HOPL: higher than optimum power load group; ES: effect size; CL: confidence limits; CMJ: countermovement jump; VEL: velocity; % chance corresponds to the probability of finding harmful/trivial/beneficial within-group changes in the performance tests.




Figure 4. Standardized mean differences for the comparisons of the between-group delta changes in the countermovement jump (CMJ) height, sprint velocities (VEL) in 5-, 10-, and 20-m, Zigzag change of direction velocity, and the relative peak power in the jump squat exercise using loads corresponding to the optimum power load (OPL), 20% lower than the OPL (-20% OPL), and 20% higher than the OPL (+20% OPL). LOPL: lower than OPL group; HOPL: higher than OPL group; the grey area represents the smallest worthwhile difference which corresponds to a small effect size (0.2); error bars represent the 90% confidence limits; #likely difference in relation to HOPL group.

composed of elastic bands) favors adaptations at the high-velocity, low-force end of the force-velocity curve. In contrast, decreasing bar-velocity (by adding traditional weights to the barbell) during JS favors adaptations at the low-velocity, high-force end of the curve (Loturco et al., 2015). Interestingly, in the current study, both training strategies were capable of enhancing power output at distinct force-velocity zones (Figure 3), which could be a direct consequence of training with load intensities near to the OPL (i.e., $\pm 20\%$ OPL). Nonetheless, the light-load group (LOPL) improved power production at all assessed zones (including at the heavier zone), whereas the heavy-load group (HOPL) increased power output only at the OPL and +20% OPL. As previously suggested, it is likely that lighter loading conditions elicit greater gains in power-related capabilities, especially when these loads are utilized in ballistic exercises (e.g., JS) (Cronin et al., 2001; Newton & Kraemer, 1994; Cormie et al., 2011b). Although the mechanisms behind this apparent superiority are unclear, it could be speculated that the higher movement velocities achieved with lighter loads may increase the rate of neural activation (by changing the pattern of motoneuron firing frequency) and provoke greater adaptations in the intermuscular coordination by, among other things, reducing the coactivation of the antagonist muscles (Cormie, McGuigan, & Newton, 2011a; Cormie et al., 2011b). These factors possibly impact the power production not only at the high-velocity zones, but across different ends of the force-velocity curve, including at the low-velocity, high-force portion. This appears to be an extra advantage in elite soccer, since light-load training probably produces lower levels of fatigue than heavy-load training, allowing players to effectively execute their technical and tactical practices (Banyard, Tufano, Delgado, Thompson, & Nosaka, 2019; Gonzalez-Badillo, Pareja-Blanco, & Rodriguez-Rosell et al., 2015; Pareja-Blanco, Rodriguez-Rosell, & Sanchez-Medina et al., 2017).

Improvements in sprinting and jumping performance are usually small (or even nonexistent) during soccer preseasons

(Loturco et al., 2017, 2015; Meckel et al., 2014; Mercer et al., 2014; Taylor et al., 2012). Loturco et al. (2015) analyzed the effects of JS or half-squat executed at the OPL throughout a 4-week preseason phase and noted that both exercises were only capable of “counteracting” the speed and power decrements in professional soccer players. Likewise, Meckel et al. (2014) observed that both continuous and interval training methods induced significant increases in aerobic fitness in young soccer players after a short-term preseason, however, these approaches also lead to stagnation or deterioration in anaerobic performance (e.g., vertical jumps). These chronic responses seem to be commonplace in various team-sport disciplines, which, as previously mentioned, may suffer negative consequences due to the interference phenomenon between concurrent aerobic and strength-power training (Docherty & Sporer, 2000; Helgerud et al., 2011; Kraemer et al., 2004). Importantly, these adverse effects can also hamper the adequate evolution and maintenance of strength, power, and speed capacities across the competitive (in-season) periods (dos Remedios, dos Remedios, & Loy et al., 1995; Segovia, Andres, Wong, & Gonzalez-Badillo, 2014; Taylor et al., 2012), which may compromise athlete performance and increase the risk of injury during matches (Lee, Mok, Chan, Yung, & Chan, 2018; van der Horst, Smits, Petersen, Goedhart, & Backx, 2015). As a consequence, the development of novel and more suitable resistance training schemes is a current and critical issue in soccer. Besides its easy implementation (the OPL can be determined by rapidly assessing bar-velocity or jump height (Loturco et al., 2015)) and apparent effectiveness (Loturco et al., 2016, 2016, 2015), the opportunity to use the OPL as a basis for defining lighter or heavier loading intensities emerges as a new strategy to enhance the functional performance of elite soccer players in different training phases (or according to the athletes’ needs). For example, our data showed that HOPL was superior for increasing COD speed and CMJ height, whereas LOPL was more efficient for improving very-short sprint performance (i.e., VEL 5-m) (Table 2). To some extent, these results are in accordance with previous studies that found meaningful improvements in COD speed in team-sport players who trained at (or close to) the OPL (Freitas, Calleja-Gonzalez, Carlos-Vivas, Marin-Cascales, & Alcaraz, 2019; Loturco et al., 2015, 2016, 2016) and greater increases in speed (e.g., 5- and 10-m) in those who executed JS at higher velocities (when compared to a “decreased velocity group”) (Loturco et al., 2015). Nevertheless, all these investigations were carried out over short periods of time (i.e., ≤ 6 weeks), making it difficult to determine the long-term effects of training under optimum loading conditions. This should certainly be addressed in future studies with longer follow-up periods.

Finally, it is important to note that we employed a restricted number of functional tests including COD, linear speed, and jump tests, which is a common and consistent practice in studies involving elite soccer players (Loturco et al., 2015, 2016, 2016). However, soccer-specific tasks (e.g., kicking, jumping to contest ball possession, tackling, etc.) may benefit from increases in the power output at distinct zones of the force-velocity curve. These technical and physical capabilities were not assessed in this research. It is probable that the OPL-based methods used here (especially the LOPL) may positively influence these critical game actions, supporting their utilization as a novel and promising training strategy for soccer athletes. This research is limited by its short duration (i.e., 4 weeks) and the

use of a single exercise (i.e., JS) in the experimental design. In contrast, the intervention was conducted throughout an actual soccer preseason, with players competing in the most important regional Brazilian youth tournament, which reinforces its applicability and ecological validity. We also recognize that (with the exception of the PP values and VEL 5-m) the majority of physical improvements detected here were “small” (ES varying from 0.23 to 0.41), which is a regular occurrence in preseason conditioning programs (Loturco et al., 2015; Meckel et al., 2014). Further studies using different exercises and more varied training approaches (e.g., combining both HOPL and LOPL regimes) are required to confirm and extend our findings. Moreover, it is recommended that the effectiveness of these training strategies be verified over long-term interventions, especially during the competitive phase of the soccer season.

Conclusion

This work has important practical implications which can be summarized as follows: 1) the OPL is possibly the heaviest loading intensity able to enhance power production under light and very-light load conditions in soccer players during congested training periods. This is reinforced by a previous study which compared the effects of OPL versus traditional strength-power periodization (Loturco et al., 2016); 2) JS training at higher loads (e.g., OPL +20%) may be necessary for improving COD performance in team-sport athletes. This conclusion is based on the current data and preliminary investigations demonstrating the importance of vertical force production in COD performance (Schreurs, Benjaminse, & Lemmink, 2017); and 3) loading ranges “immediately” below the OPL (i.e., OPL –20%) appear to be effective for increasing very-short sprint ability (i.e., 5-m) in soccer players, even during short preseasons. A probable explanation for this effectiveness is related to the lower levels of fatigue generated by light loads (Gonzalez-Badillo et al., 2015), which is certainly a great advantage in elite soccer settings (especially when considering the critical role of maximum acceleration and speed in modern soccer) (Barnes, Archer, Hogg, Bush, & Bradley, 2014; Faude et al., 2012). Soccer coaches and sport scientists can implement the JS OPL-based training schemes presented here, either separately or combined, according to individual necessities and specific playing tasks.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Irineu Loturco  <http://orcid.org/0000-0003-1309-2568>
 Pedro E. Alcaraz  <http://orcid.org/0000-0002-9792-6656>
 Tomás T. Freitas  <http://orcid.org/0000-0001-8571-3189>

References

Banyard, H. G., Tufano, J. J., Delgado, J., Thompson, S. W., & Nosaka, K. (2019). Comparison of the effects of velocity-based training methods and traditional 1RM-percent-based training prescription on acute

- kinetic and kinematic variables. *International Journal of Sports Physiology and Performance*, 14, 246–255.
- Barnes, C., Archer, D. T., Hogg, B., Bush, M., & Bradley, P. S. (2014). The evolution of physical and technical performance parameters in the English premier league. *International Journal of Sports Medicine*, 35, 1095–1100.
- Batterham, A. M., & Hopkins, W. G. (2006). Making meaningful inferences about magnitudes. *International Journal of Sports Physiology and Performance*, 1, 50–57.
- Behm, D. G., & Sale, D. G. (1993). Velocity specificity of resistance training. *Sports Medicine*, 15, 374–388.
- Brown, L., & Whitehurst, M. (2003). The effect of short-term isokinetic training on force and rate of velocity development. *Journal of Strength and Conditioning Research*, 17, 88–94.
- Cormie, P., McGuigan, M. R., & Newton, R. U. (2011a). Developing maximal neuromuscular power: Part 1-biological basis of maximal power production. *Sports Medicine*, 41, 17–38.
- Cormie, P., McGuigan, M. R., & Newton, R. U. (2011b). Developing maximal neuromuscular power: Part 2 - training considerations for improving maximal power production. *Sports Medicine*, 41, 125–146.
- Coyle, E. F., Feiring, D. C., Rotkis, T. C., Cote 3rd, R. W., Roby, F. B., Lee, W., & Wilmore, J. H. (1981). Specificity of power improvements through slow and fast isokinetic training. *Journal of Applied Physiology: Respiratory, Environmental and Exercise Physiology*, 51, 1437–1442.
- Cronin, J., McNair, P. J., & Marshall, R. N. (2001). Developing explosive power: A comparison of technique and training. *Journal of Science and Medicine in Sport*, 4, 59–70.
- Docherty, D., & Sporer, B. (2000). A proposed model for examining the interference phenomenon between concurrent aerobic and strength training. *Sports Medicine*, 30, 385–394.
- dos Remedios, K. A., dos Remedios, R. L., Loy, S. F., Holland, G. J., Vincent, W. J., Conley, L. M., & Heng, M. (1995). Physiological and field test performance changes of community college football players over a season. *Journal of Strength and Conditioning Research*, 9, 211–215.
- Faude, O., Koch, T., & Meyer, T. (2012). Straight sprinting is the most frequent action in goal situations in professional football. *Journal of Sports Sciences*, 30, 625–631.
- Freitas, T. T., Calleja-Gonzalez, J., Carlos-Vivas, J., Marin-Cascales, E., & Alcaraz, P. E. (2019). Short-term optimal load training vs a modified complex training in semi-professional basketball players. *Journal of Sports Sciences*, 37, 434–442.
- Gonzalez-Badillo, J. J., Pareja-Blanco, F., Rodriguez-Rosell, D., Abad-Herencia, J. L., del Ojo-López, J. J., & Sánchez-Medina, L. (2015). Effects of velocity-based resistance training on young soccer players of different ages. *Journal of Strength and Conditioning Research*, 29, 1329–1338.
- Helgerud, J., Rodas, G., Kemi, O. J., & Hoff, J. (2011). Strength and endurance in elite football players. *International Journal of Sports Medicine*, 32, 677–682.
- Hopkins, W. G., Marshall, S. W., Batterham, A. M., & Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise*, 41, 3–13.
- Kanehisa, H., & Miyashita, M. (1983). Specificity of velocity in strength training. *European Journal of Applied Physiology*, 52, 104–106.
- Kawamori, N., & Haff, G. G. (2004). The optimal training load for the development of muscular power. *Journal of Strength and Conditioning Research*, 18, 675–684.
- Kraemer, W. J., French, D. N., Paxton, N. J., Hakkinen, K., Volek, J. S., Sebastianelli, W. J., ... Knuttgen, H. G. (2004). Changes in exercise performance and hormonal concentrations over a big ten soccer season in starters and nonstarters. *Journal of Strength and Conditioning Research*, 18, 121–128.
- Lee, J. W. Y., Mok, K. M., Chan, H. C. K., Yung, P. S. H., & Chan, K. M. (2018). Eccentric hamstring strength deficit and poor hamstring-to-quadriceps ratio are risk factors for hamstring strain injury in football: A prospective study of 146 professional players. *Journal of Science and Medicine in Sport*, 21, 789–793.
- Loturco, I., Kobal, R., Kitamura, K., Cal Abad, C. C., Faust, B., Almeida, L., & Pereira, L. A. (2017). Mixed training methods: Effects of combining resisted sprints or plyometrics with optimum power loads on sprint

- and agility performance in professional soccer players. *Frontiers in Physiology*, 8, 1034.
- Loturco, I., Kobal, R., Moraes, J. E., Kitamura, K., Cal Abad, C. C., Pereira, L. A., & Nakamura, F. Y. (2017). Predicting the maximum dynamic strength in bench press: The high precision of the bar velocity approach. *Journal of Strength and Conditioning Research*, 31, 1127–1131.
- Loturco, I., Nakamura, F. Y., Kobal, R., Gil, S., Cal Abad, C. C., Cuniyochi, R., ... Roschel, H. (2015). Training for power and speed: Effects of increasing or decreasing jump squat velocity in elite young soccer players. *Journal of Strength and Conditioning Research*, 29, 2771–2779.
- Loturco, I., Nakamura, F. Y., Kobal, R., Gil, S., Pivetti, B., Pereira, L. A., & Roschel, H. (2016). Traditional periodization versus optimum training load applied to soccer players: Effects on neuromuscular abilities. *International Journal of Sports Medicine*, 37, 1051–1059.
- Loturco, I., Nakamura, F. Y., Tricoli, V., Kobal, R., Abad, C. C., Kitamura, K., ... Gonzalez-Badillo, J. J. (2015). Determining the optimum power load in jump squats using the mean propulsive velocity. *PLoS One*, 10, e0140102.
- Loturco, I., Pereira, L. A., Kobal, R., Zanetti, V., Gil, S., Kitamura, K., ... Nakamura, F. Y. (2015). Half-squat or jump squat training under optimum power load conditions to counteract power and speed decrements in Brazilian elite soccer players during the preseason. *Journal of Sports Sciences*, 33, 1283–1292.
- Loturco, I., Pereira, L. A., Kobal, R., Maldonado, T., Piazza, A. F., Bottino, A., ... Nakamura, F. Y. (2016). Improving sprint performance in soccer: Effectiveness of jump squat and Olympic push press exercises. *PLoS One*, 11, e0153958.
- Loturco, I., Pereira, L. A., Kobal, R., & McGuigan, M. R. (2019). Power output in traditional and ballistic bench press in elite athletes: Influence of training background. *Journal of Sports Sciences*, 37, 277–284.
- McBride, J. M., Triplett-McBride, T., Davie, A., & Newton, R. U. (2002). The effect of heavy- vs. light-load jump squats on the development of strength, power, and speed. *Journal of Strength and Conditioning Research*, 16, 75–82.
- Meckel, Y., Harel, U., Michaely, Y., & Eliakim, A. (2014). Effects of a very short-term preseason training procedure on the fitness of soccer players. *The Journal of Sports Medicine and Physical Fitness*, 54, 432–440.
- Mercer, T. H., Gleeson, N. P., & Mitchell, J. (2014). *Fitness profiles of professional soccer players before and after a preseason conditioning*. Abingdon: Routledge.
- Newton, R. U., & Kraemer, W. J. (1994). Developing explosive muscular power: Implications for a mixed methods training strategy. *Strength and Conditioning Journal*, 16, 20–31.
- Noon, M. R., James, R. S., Clarke, N. D., Akubat, I., & Thake, C. D. (2015). Perceptions of well-being and physical performance in English elite youth footballers across a season. *Journal of Sports Sciences*, 18, 1–10.
- Pareja-Blanco, F., Rodriguez-Rosell, D., Sanchez-Medina, L., Sanchis-Moysi, J., Dorado, C., Mora-Custodio, R., ... Gonzalez-Badillo, J. J. (2017). Effects of velocity loss during resistance training on athletic performance, strength gains and muscle adaptations. *Scandinavian Journal of Medicine & Science in Sports*, 27, 724–735.
- Sanchez-Medina, L., Perez, C. E., & Gonzalez-Badillo, J. J. (2010). Importance of the propulsive phase in strength assessment. *International Journal of Sports Medicine*, 31, 123–129.
- Schreurs, M. J., Benjaminse, A., & Lemmink, K. (2017). Sharper angle, higher risk? The effect of cutting angle on knee mechanics in invasion sport athletes. *Journal of Biomechanics*, 63, 144–150.
- Segovia, M. L., Andres, J. M. P., Wong, D. P., & Gonzalez-Badillo, J. J. (2014). Changes in strength and aerobic performance by concurrent training in under-19 soccer players. *International SportMed Journal*, 15, 123–135.
- Taylor, J. M. P., Portas, M. D., Wright, C. H., & Weston, M. (2012). Within-season variation of fitness in elite youth female soccer players. *Journal of Athletic Enhancement*, 1, 1–5.
- Tufano, J. J., Brown, L. E., & Haff, G. G. (2017). Theoretical and practical aspects of different cluster set structures: A systematic review. *Journal of Strength and Conditioning Research*, 31, 848–867.
- van der Horst, N., Smits, D. W., Petersen, J., Goedhart, E. A., & Backx, F. J. (2015). The preventive effect of the nordic hamstring exercise on hamstring injuries in amateur soccer players: A randomized controlled trial. *The American Journal of Sports Medicine*, 43, 1316–1323.