SHORT-TERM TRAINING AND DETRAINING EFFECTS OF SUPERVISED VS. UNSUPERVISED RESISTANCE EXERCISE IN AGING ADULTS

SAMUEL T. ORANGE,¹ PHIL MARSHALL,¹ LEIGH A. MADDEN,² AND REBECCA V. VINCE¹

¹Sport, Health and Exercise Science, School of Life Sciences, University of Hull, Hull, United Kingdom; and ²Center of Biomedical Research, School of Life Sciences, University of Hull, Hull, United Kingdom

Abstract

Orange, ST, Marshall, P, Madden, LA, and Vince, RV. Shortterm training and detraining effects of supervised vs. unsupervised resistance exercise in aging adults. J Strength Cond Res 33(10): 2733-2742, 2019-This study compared the effects of a 4-week supervised (SUP) resistance training program and unsupervised (UNSUP) resistance training program followed by 12 weeks of detraining (DET). Thirty-six healthy aging adults (age: 53.6 \pm 3.6 years; body mass index: 28.3 \pm 5.1 kg·m⁻²) were randomly allocated to an SUP group (n = 17) or a UNSUP group (n = 19). Participants completed 3 training sessions per week using resistance bands and body weight movements. Measures of physical performance were administered at baseline, at the end of the training program, and after the DET period. Function was assessed with the 6-minute walk test (6MWT), timed up-and-go (TUG), 30-second chair sit-to-stand (STS), stair-climb test (SCT), 40-m fast-paced walk test (FPWT) and sit-and-reach test (SRT), whereas the isometric midthigh pull (IMTP) and hand grip test were used to measure muscle strength. After training, improvements in performance were found in the 6MWT, TUG, 30-second chair STS, SCT, FPWT, SRT, and IMTP ($p \le 0.05$), with no significant differences between groups (p > 0.05). In addition, most of the training-induced improvements remained significantly above baseline values after the DET period ($p \leq 0.05$). No significant between-group differences were observed after training or DET (p > 0.05). Four weeks of either SUP or UNSUP resistance training is sufficient to substantially improve muscle strength and function in aging adults, and these gains are largely preserved after prescribed exercise cessation.

33(10)/2733-2742

Journal of Strength and Conditioning Research © 2018 National Strength and Conditioning Association Home-based resistance training seems to be a practical and effective alternative to traditional SUP programs that may help circumvent many barriers to physical activity in aging adults.

KEY WORDS resistance training, functional capacity, homebased exercise, body weight exercise, resistance band exercise

INTRODUCTION

egular exercise opposes the debilitating effects of aging by mitigating declines in muscle strength and function (42). In particular, progressive resistance training has consistently been shown to improve functional abilities in adults aged 50 years or older (18,21,38). However, the research area is currently dominated by gym-based interventions requiring specialized equipment and personnel, with little consideration for long-term sustainability (11). A lack of access to transportation and traditional resistance facilities limits the widespread application of resistance training to this discrete population. In fact, older individuals are more likely to engage in exercise interventions that are easily accessible, do not require transport, and involve no out-of-pocket costs (23).

Home-based exercise is a convenient alternative to supervised (SUP) programs and may promote greater long-term participation than exercising at a designated setting (3). Despite the clear economic and practical benefits of home-based exercise, a recent systematic review (37) has suggested that SUP resistance training improves measures of muscle strength to a greater extent than unsupervised (UNSUP) programs. It is pertinent to note, however, that the limited work systematically comparing these 2 intervention strategies have used SUP group exercise sessions, whereas the UNSUP home exercise has been performed individually. Given that social interaction is a robust and well-established exercise motive for older adults (20), it is conceivable that these comparisons were confounded by the social element of group training. Delivering both interventions on an individual basis would better identify the impact that supervision alone has on exercise-derived functional benefits in aging adults. It is also important that UNSUP resistance training programs still pay attention to

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (http://journals.lww.com/nsca-jscr).

Address correspondence to Phil Marshall, phil.marshall@hull.ac.uk.

the fundamental principles of exercise physiology to strike a balance between efficacious and sustainable training.

Perhaps, the hallmark of an effective resistance training program, the principle of specificity asserts that the training stimulus must be specific to the desired adaptation (41). That is, the exercise must replicate the biomechanical movement patterns and underpinning bioenergetics involved in the performance of the primary outcome measurement (27). To improve function, training movements should promote the transfer of force through lower-body triple extension to simulate activities of everyday life, such as rising from a chair and climbing the stairs. It is therefore surprising that most of the training studies in older persons have primarily involved single-joint movements and the use of resistance machines (18,21,25). Training with elastic bands enables the execution of functional movement patterns through a full range of motion (ROM). Moreover, multiarticular exercises can easily be performed in multiple planes of motion because the direction of the resistance depends on the positioning of the elastic device rather than gravity (32). Importantly, elastic bands and free weights have been shown to exert similar benefits on measures of functional capacity in older adults (17).

The extent to which training-induced adaptations can be maintained after the cessation of prescribed exercise is a necessary consideration for any training program. Studies have shown periods of detraining (DET) to retain (31,36,48) or completely reverse (18) measures of musculoskeletal strength and function after systematized resistance training. It is currently unknown whether supervision mediates the effect that DET has on physical performance in aging individuals. Therefore, the aims of this study were threefold: (a) to examine the effectiveness of a short-term functional resistance training program on measures of muscle strength and function in adults aged 50–65 years, (b) to compare the efficacy of resistance training performed in an SUP vs. UNSUP setting, and (c) to determine whether changes in muscle strength and function are maintained after a DET period of 12 weeks. Based on the current literature, we hypothesized that (a) resistance training would result in significant improvements in all outcome measurements, (b) the SUP group would improve strength and functional performance to a greater extent than the UNSUP group, and (c) training-induced improvements would remain above baseline levels after the DET period in both training groups.

METHODS

Experimental Approach to the Problem

This study was a 2-arm experimental trial whereby participants were randomly allocated to an SUP or UNSUP group. Both groups completed 4 weeks of functional resistance training with all variables controlled between conditions apart from the level of supervision. Outcomes measurements of functional and physical performance were administered at baseline (before group allocation), at the end of the 4-week intervention, and after a DET period of 12 weeks. All participants agreed to maintain their current diet and activity levels during the intervention period. There were no particular instructions or guidance given during the 12-week DET phase.

TABLE 1. Baseline characteristics of study participants.*†							
	Total (<i>n</i> = 36)	SUP (<i>n</i> = 17)	UNSUP $(n = 19)$	р			
Age (y)	$53.6~\pm~3.6$	52.9 ± 3.8	$54.2~\pm~3.3$	0.295			
Males/females	11/25	4/13	7/12	0.510			
Body mass (kg)	78.0 ± 16.5	76.1 ± 17.4	79.7 ± 15.9	0.526			
Height (cm)	165.9 \pm 9.5	164.1 ± 9.5	167.5 ± 9.4	0.283			
BMI (kg⋅m ⁻²)	$\textbf{28.3}~\pm~\textbf{5.1}$	$\textbf{28.1}~\pm~\textbf{4.9}$	$28.4~\pm~5.5$	0.832			
Blood pressure							
Systolic (mm Hg)	132.3 ± 11.4	131.4 ± 12.6	133.1 ± 10.6	0.653			
Diastolic (mm Hg)	$84.7~\pm~8.5$	$83.8~\pm~9.6$	$85.5~\pm~7.6$	0.557			
Resting HR (b ⋅ min ⁻¹)	$72.7~\pm~7.3$	$72.4~\pm~8.3$	$73.0~\pm~6.5$	0.813			
6MWT (m)	614.79 ± 53.33	614.23 ± 59.79	615.28 ± 48.50	0.954			
TUG (s)	6.39 ± 0.65	6.33 ± 0.62	$6.44~\pm~0.68$	0.592			
30-s chair STS (reps)	$12.8~\pm~2.2$	13.1 ± 2.2	$12.6~\pm~2.2$	0.517			
SCT (s)	5.86 ± 0.78	5.80 ± 0.83	5.91 ± 0.76	0.676			
FPWT (s)	20.46 ± 1.76	20.51 ± 2.06	20.41 ± 1.50	0.871			
Hand grip (kg)	34.8 ± 10.0	$33.0~\pm~8.3$	36.5 ± 11.3	0.302			
IMTP (kg)	$79.4~\pm~39.6$	78.4 ± 35.7	$80.4~\pm~43.7$	0.881			
SRT (cm)	15.8 ± 9.7	$17.4~\pm~9.6$	14.3 ± 9.7	0.332			

*SUP = supervised; UNSUP = unsupervised; BMI = body mass index; HR = heart rate; 6MWT = 6-minute walk test; TUG = timed up-and-go; STS = sit-to-stand; SCT = stair-climb test; FPWT = fast-paced walk test; IMTP = isometric midthigh pull; SRT = sit-and-reach test. †Data are presented as mean \pm SD.

2734 Journal of Strength and Conditioning Research

Subjects

All participants were required to be aged 50-65 years, have a body mass index of less than 35 kg·m⁻², have not engaged in more than 30 minutes of moderate-vigorous intensity exercise on 3 or more days of the week for the past 3 months, and have no resistance training experience in the past 12 months. In total, 36 participants completed the resistance training intervention, with 17 in the SUP group and 19 in the UNSUP group (see Figure, Supplemental Digital Content 1, http://links. lww.com/JSCR/A79). Participants were informed of the experimental procedures to be undertaken before signing an institutionally approved informed consent document to participate in the study. Baseline characteristics of study

participants are presented in Table 1. The study was approved by the Sport, Health and Exercise Science Ethics Committee at the University of Hull.

Procedures

Resistance Training Program. The resistance training program was designed and delivered by a Certified Strength and Conditioning Specialist (CSCS) and was based on guidelines by the National Strength and Conditioning Association (NSCA) (27). Participants in both groups completed 3 sessions per week on nonconsecutive days for the 4-week intervention period. One set of 8 repetitions was performed in week 1, 2 sets of 8 repetitions in week 2, 2 sets of 10 repetitions in week 3, and 3 sets of 10 repetitions in week 4. The intensity of exercise was performed at 4-6 on the modified 10-point rating of perceived exertion (RPE) scale (13) associated with the target number of repetitions. This corresponded with qualitative descriptions of "somewhat hard" to "hard." After a dynamic warm-up that included targeted mobility exercises designed to increase the ROM in the ankle, hip, and thoracic spine, participants performed 11 resistance exercises using body weight and resistance bands (Iron Woody Fitness, Olney, MT, USA). Each exercise was based on a primary resistance training movement pattern as described in Table 2. Three color-coded bands were used offering 3 incremental levels of resistance (yellow, purple and red for light, medium, and heavy resistance, respectively). Each session finished with a cool-down of static stretching that included ankle, hip, gluteal, hamstring, and pectoral stretches.

The exercises focused on multiarticular and multiplanar movements to provide a functional training stimulus and mimic activities of daily living. Exercises were sequenced so that upper- and lower-body movements were alternated, which has been suggested to be beneficial for untrained individuals who may find that completing several lower- or upper-body exercises in succession is too strenuous (27). Progression (and regression) of the training load and volume was based on the participant's RPE rating. If RPE was below 4 or above 6, the exercise was progressed or regressed for the next workout, respectively. The resistance band exercises were progressed by changing from the current band to the next color in the scale (e.g., yellow to purple). If a participant reached the level of resistance red, another band was added while following the progression scale (e.g., red plus yellow). Body weight exercises were progressed using exercises of similar movement patterns with a higher degree of technical difficulty (e.g., biped stance to split stance).

Unsupervised Training Program. Participants completed the UNSUP training program individually in their home. After completion of the baseline assessments, participants returned to the laboratory to be familiarized with the exercises to be used in the study and the use of the modified RPE scale. A CSCS checked for correct form in all exercises and adjusted technique if necessary. Participants then received an exercise package that included 3 color-coded resistance bands, a heart rate monitor (FT1; Polar Electro, Kempele, Finland), an exercise DVD, a training log, an exercise progression/ regression sheet, and the modified RPE Scale. An instructional booklet was also included, written in layman language with images and diagrams, clearly describing all components of the program. The CSCS telephoned all participants once per week to answer any questions and to document their RPE rating for each exercise. If participants' RPE for a given exercise fell outside the predetermined level of intensity

Movement	Key exercise	
Hip extension†	Shoulder-raised bilateral glute bridge	
Lower-body triple extension†	Squat	
Horizontal push [†]	Modified press-up	
Lower-body triple extension†	Split squat	
Scapula retraction [‡]	Standing scapula retraction w/yellow band	
Lateral rotatory	Lateral walk w/yellow band	
Vertical pusht	Incline chest press w/yellow band	
Hip hinget	Deadlift w/yellow band	
Horizontal pull‡	Seated row w/yellow band	
Full-body extension [‡]	Push press w/yellow band	
Anti-rotation [‡]	Core rotation w/yellow band	

*The resistance exercises were based on primary resistance training movement patterns. Key exercises used in the intervention are shown here. These key exercises were modified according to the participants' rating of perceived exertion. †Body weight exercise.

 \ddagger Resistance band exercise; w/ = with.

(RPE of below 4 or above 6), they were prompted to use their exercise progression/ regression sheet to modify the exercise accordingly.

Supervised Training Program. The SUP group followed the same exercise program as the UNSUP group, apart from that they completed the sessions in our Biomechanics laboratory and received one-to-one supervision by the same CSCS who provided telephone support to the UNSUP group. Participants received real-time encouragement and feedback on exercise technique with form being adjusted by the CSCS if necessary. Rating of perceived exertion data were

VOLUME 33 | NUMBER 10 | OCTOBER 2019 | 2735

collected after the cessation of each training session and exercises were modified for the next workout accordingly.

Outcome Measurements: Six Minute-Walk Test. Participants were instructed to walk at their own maximal pace back and forth along a flat 30-m surface, covering as much ground as they could in 6 minutes. All instructions, encouragement, and monitoring adhered to the guidelines provided by the American Thoracic Society (4). Participants completed one trial and the distance covered was recorded in meters. The 6-minute walk test (6MWT) has recently demonstrated excellent reliability in our laboratory (intraclass correlation [ICC] = 0.98), with the SEM and minimum detectable change at 95% confidence intervals (CIs) (MDC_{95%}) reported at 13.7 and 37.8 m, respectively (40).

Timed Up-and-Go. Participants sat in a firm, armless chair (height, 40 cm; depth, 39 cm) and were instructed to stand up, walk 3 m before turning 180°, and return to the chair to sit down. Participants were instructed to perform the test as quickly as possible but in a controlled manner, with time recorded in seconds during one trial. Timed up-and-go (TUG) is a basic measure of functional mobility (7) and has demonstrated high test-retest reliability in our laboratory (ICC = 0.97; SEM = 0.22 seconds; MDC_{95%} = 0.62 seconds) (40).

Thirty-Second Chair Sit-To-Stand. The 30-second chair sit-tostand (STS) is a reliable measure of lower-extremity function and strength in older adults (ICC = 0.89) (33). The test was administered using the same chair as the TUG, which was supported against a wall. Participants began seated and were subsequently instructed to rise to a full standing position (legs straight) and then return to the seat (full weight on chair) with both arms crossed against the chest. A practice trial of 2 repetitions was given to check correct form, followed by one test trial. The total number of stands performed correctly in 30 seconds was recorded for analysis.

Stair-Climb Test. Participants ascended and descended a freestanding flight of 5 steps (step height, 20 cm) as quickly possible, but in a safe and controlled manner. The use of the handrails was permitted if required, and the test finished when both feet were flat on the ground level. One trial was permitted with the time recorded in seconds. Using our laboratory's custom-built staircase, the stair-climb test (SCT) has been shown to be highly reliable (ICC = 0.98; *SEM* = 0.08 seconds; MDC_{95%} = 0.22 seconds) (40).

Forty-Meter Fast-Paced Walk Test. Participants walked as quickly as possible along a 20-m flat surface, turned 180° around a cone, and then walked 20 m back to the start line. The test finished when the participant had walked 40 m to cross back over the start line, with time recorded in seconds during one trial. The 40-m fast-paced walk test (FPWT) has previously demonstrated excellent reliability (ICC = 0.95, SEM = 1.0 m s⁻¹) (51).



 \leq 0.05). Data are presented as mean \pm SE.



Hand Grip Test. Using their dominant hand, participants squeezed the analogue dynamometer (T.K.K. 5001 Grip-A; Takei Scientific Instruments, Niigata City, Japan) as hard as possible for 2-3 seconds. An upright biped position was maintained throughout the test with the arm in full extension. The grip position of the dynamometer was adjusted to each individual's hand size. The best score of 2 trials was recorded to the nearest 0.5 kg and used for analysis. The TKK dynamometer has recently recorded high reliability and criterion-related validity (12).

Isometric Midthigh Pull. Using an analogue back dynamometer (T.K.K. 5002 Back-A; Takei Scientific Instruments, Niigata City, Japan), participants maximally extended their knees and trunk for 5 seconds without bending their back. The height of the handle was individually adjusted so that the bar rested midway up the thigh and there was 145° of knee flexion (22). Two trials were performed with a 2-minute rest period in between. Each trial was recorded to the nearest 1 kg, with the mean value used for analysis. This test has previously demonstrated good to acceptable reliability (ICC = 0.81-0.85) (26).

Sit-and-Reach Test. Sit-and-reach test (SRT) is a reliable (ICC = 0.94) (10) measure of hamstring and spinal flexibility. Participants sat on the floor with their legs fully extended and heels flat against a standardized box (height, 32.5 cm). One hand was placed on top of the other and participants gradually reached forward as far as possible along the measuring tape on top of the box. One trial was completed, and the furthest the participants reached and held for 2 seconds was recorded to the nearest 0.5 cm.

Heart Rate. Average and maximum heart rate and session duration were recorded for each training session using the Polar heart rate monitor. Recording commenced before the

start of the warm-up and stopped immediately after the last resistance exercise (before the cool-down).

Exercise Compliance. Compliance in the training intervention was calculated as follows: ([sessions attended/total number of sessions] \times 100). Participation in the SUP intervention was assessed using attendance at the SUP training sessions. Participation in the UNSUP intervention was evaluated using participants' training logs.

Sample Size Estimation. The sample size was calculated using G*Power software (version 3.1; Universität Düsseldorf, Düsseldorf, Germany). Given the type of study design (mixed analysis of variance [ANOVA] with repeated measures), the following input parameters were entered to obtain medium-sized group \times time interaction effects: $\alpha = 0.05$, statistical power of 0.8, and an effect size of 0.25. Thus, a priori sample size for statistical significance was calculated as 28 participants (i.e., 14 in each group). A dropout rate of 20% was also considered. The medium effect size was based on a recent

Outcome	PRE-POST	PRE-DET	POST-DET	Time $ imes$ group interaction p
6MWT (m)				0.849
SUP	48.6 (28.4 to 68.8)‡	30.6 (6.2 to 55.0)‡	-18.0 (-38.3 to 2.2)	
UNSUP	42.7 (17.8 to 67.7)‡	24.7 (1.6 to 47.8)‡	-18.1 (-37.8 to 1.7)	
TUG (s)				0.746
SUP	-0.84 (-1.16 to -0.53)‡	-0.61 (-0.88 to -0.34)	0.23 (-0.16 to 0.62)	
UNSUP	-0.93 (-1.28 to -0.58)‡	-0.57 (-0.88 to -0.27)	0.36 (0.13 to 0.58)‡	
30-s chair ST	S			0.784
(s)				
SUP	3.4 (2.8 to 3.9)‡	1.8 (1.0 to 2.6)‡	−1.6 (−2.6 to −0.6)‡	
UNSUP	3.1 (2.0 to 4.1)‡	1.8 (0.8 to 2.9)‡	−1.3 (−2.4 to −0.1)‡	
SCT (s)				0.923
SUP	-0.74 (-1.12 to -0.36)‡	-0.42 (-0.78 to -0.05)	0.32 (0.07 to 0.57)‡	
UNSUP	-0.79 (-1.17 to -0.41)‡	-0.44 (-0.78 to -0.11)	0.34 (0.14 to 0.55)‡	
FPWT (s)				0.299
SUP	-1.74 (-2.87 to -0.62)‡	-0.93 (-1.42 to -0.44)	∴ 0.81 (−0.48 to 2.10)	
UNSUP	-1.37 (-1.98 to -0.76)‡	-1.22 (-1.85 to -0.60)	0.15 (-0.28 to 0.57)	
Hand grip te	st			0.140
(kg)				
SUP	0.9 (-1.0 to 2.7)	1.6 (−0.3 to 3.5)	0.7 (-0.7 to 2.2)	
UNSUP	0.8 (-0.2 to 1.8)	0.3 (-0.7 to 1.2)	-0.6 (-1.8 to 0.7)	
MTP (kg)				0.829
SUP	25.0 (16.4 to 33.6)‡	14.4 (7.2 to 21.6)‡	−10.6 (−18.3 to −3.0)‡	
UNSUP	26.6 (14.3 to 38.8)‡	17.6 (4.8 to 30.5)‡	−8.9 (−17.6 to −0.3)‡	
SRT (cm)				0.495
SUP	3.2 (−0.2 to 6.6)‡	2.1 (-1.7 to 6.0)	-1.0 (-3.2 to 1.1)	
UNSUP	3.2 (1.3 to 5.1)‡	0.9 (-0.8 to 2.6)	−2.3 (−3.7 to −1.0)‡	

*PRE = preintervention; POST = postintervention; DET = detraining; 6MWT = 6-minute walk test; SUP = supervised; UNSUP = unsupervised; TUG = timed up-and-go; STS = sit-to-stand; SCT = stair-climb test; FPWT = fast-paced walk test; IMTP = isometric midthigh pull; SRT = sit-and-reach test.

†Data are presented as mean values (95% confidence intervals).

 \pm Significant difference within groups ($p \le 0.05$).

TABLE 4. Between-group changes between the different time points.*†						
	PRE-POST	PRE-DET	POST-DET			
6MWT (m) TUG (s) 30-s chair STS (reps) SCT (s) FPWT (s) Hand grip test (kg) IMTP (kg) SRT (cm)	5.89 (-19.08 to 30.86) 0.09 (-0.28 to 0.45) 0.30 (-0.64 to 1.24) 0.05 (-0.36 to 0.47) 0.37 (-0.58 to 1.32) 0.04 (-1.52 to 1.59) 1.58 (-10.17 to 13.32) 0.03 (-2.88 to 2.95)	5.93 (-19.78 to 31.65) 0.04 (-0.28 to 0.36) 0.03 (-1.03 to 1.08) 0.28 (-0.35 to 0.40) 0.29 (-0.33 to 0.91) 1.33 (-0.23 to 2.88) 3.3 (-8.41 to 14.97) 1.28 (-1.84 to 4.39)	$\begin{array}{c} 0.41 \ (-21.65 \ {\rm to} \ 21.73) \\ 0.13 \ (-0.21 \ {\rm to} \ 0.46) \\ 0.33 \ (-0.83 \ {\rm to} \ 1.48) \\ 0.24 \ (-0.22 \ {\rm to} \ 0.27) \\ 0.66 \ (-0.33 \ {\rm to} \ 1.7) \\ 1.29 \ (-0.18 \ {\rm to} \ 2.75) \\ 1.7 \ (-7.2 \ {\rm to} \ 10.61) \\ 1.31 \ (-0.56 \ {\rm to} \ 3.19) \end{array}$			

*PRE = preintervention; POST = postintervention; DET = detraining; 6MWT = 6-minute walk test; TUG = timed up-and-go; STS = sit-to-stand; SCT = stair-climb test; FPWT = fast-paced walk test; IMTP = isometric midthigh pull; SRT = sit-and-reach test. †Data are presented as mean values (95% confidence interval).

meta-analysis (37) comparing the effects of SUP vs. UNSUP resistance training on measures of muscle strength in older adults (standardized mean difference [SMD] = 0.51).

Statistical Analyses

All analyses were performed by intention to treat using SPSS for Windows (IBM SPSS, version 22.0, Chicago, IL, USA). Shapiro-Wilk and Levene's tests were used to verify normality of data and homogeneity of variance, respectively, and all assumptions were met. To compare baseline characteristics between groups, an independent-samples t-test was conducted for continuous variables, whereas the Mann-Whitney U-test was used for ordinal data (sex). A 2 \times 3 mixed-model ANOVA with repeated measures for group (between) and time (within) was used to examine the effects of the intervention on each outcome measurement. The alpha level indicating statistical significance for this test was set at $p \leq 0.05$. The data were then further explored with pairwise comparisons using a Bonferroni-adjusted alpha level. The assumption of sphericity was assessed with Mauchly's test, and in the case of significant violations, the Greenhouse-Geisser epsilon correction was applied. The level for all CIs was 95%.

RESULTS

Exercise Responses

Exercise compliance was 94.6% in the SUP group and 98.7% in the UNSUP group, with no significant difference between conditions (4.1 \pm 2.1%, p = 0.066, 95% CI: -8.4 to 0.3%). Session duration was 27.6 \pm 2.9 minutes in the SUP group and 23.1 \pm 3.4 minutes in the UNSUP group, with this difference reaching statistical significance (5 \pm 1 minute, $p \leq 0.05$, 95% CI: 2–7 minutes). There was a significant interaction between group and time on average heart rate ($p \leq 0.05$). Specifically, average heart rate was 14 \pm 3 b min⁻¹ higher in the SUP group compared with the UNSUP group ($p \leq 0.05$, 95% CI: 7–22 b min⁻¹) (Figure 1). For maximum

heart rate, there was no significant group by time interaction (p = 0.770). However, there were significant main effects of time ($p \le 0.05$) and group ($p \le 0.05$), showing that peak heart rate was 23 ± 4 b·min⁻¹ ($p \le 0.05$; 95% CI: 14–33 b·min⁻¹) higher in the SUP group compared with the UNSUP group.

Physical Performance Outcomes

There were no significant main effects of group nor any significant interaction effects between group and time for any physical performance outcome (p > 0.05; Table 3). However, the main effect of time showed a statistically significant difference in all variables at the different time points $(p \le 0.05)$. With the exception of hand grip strength, Bonferroni-corrected pairwise comparisons revealed significant training-induced improvements in all performance tasks $(p \le 0.05;$ Table 3). Detraining resulted in significant reductions in 30-second chair STS, SCT, and isometric midthigh pull (IMTP) performance in both the SUP and UNSUP conditions ($p \le 0.05$; Table 3). Timed up-and-go and SRT performance also significantly decreased in the UNSUP group but not in the SUP group after DET, although these reductions were not significantly different between conditions (TUG: 0.13 seconds, p = 0.454, 95% CI: -0.21 to 0.46 seconds; SRT: 1.31 cm, p = 0.924, 95% CI: -0.56 to 3.19 cm). Despite these performance decrements, the 6MWT, TUG, 30-second chair STS, SCT, FPWT, and IMTP remained significantly above baseline in both groups after DET ($p \le 0.05$). No significant between-group differences emerged between conditions after training or DET (p > 0.05) (Table 4).

DISCUSSION

This study examined the short-term training and DET effects of SUP vs. UNSUP resistance training on muscle strength and function in adults aged 53.6 \pm 3.6 years. Our data demonstrate a comparative increase in functional ability and muscle strength after both training interventions. These

Copyright © 2018 National Strength and Conditioning Association. Unauthorized reproduction of this article is prohibited.

improvements were attained using low-cost elastic bands and a small weekly time commitment (83 and 69 minutes in SUP and UNSUP, respectively). In addition, most of the training-induced adaptations remained above baseline values after the period of DET.

The training program resulted in a significant increase in functional performance and IMTP strength, independent of the level of supervision. The magnitude of change, considered in relation to the error of measurement, suggests that the training-induced improvements are likely to be meaningful for aging adults. For example, the improved TUG performance observed in both groups (-0.84 and -0.93)seconds in SUP and UNSUP, respectively) exceeds the SEM (0.22 seconds) and MDC_{95%} (0.62 seconds) previously recorded in our laboratory (40). This improvement in functional performance is also greater than the magnitude reported in a previous meta-analysis of resistance training in older adults (-0.69 seconds, 95% CI: -1.11 to -0.27 seconds) (38) and is larger than the change observed in a number of recent studies in this area (16,47). This difference may be attributed to the average age of participants. In this study, subjects had a mean age of 53.6 \pm 3.6 years, whereas the mean age of trials included in the meta-analytic review (38)ranged from 65.8 ± 7.6 to 84.9 ± 4.8 years. Alternatively, the difference in magnitude may be related to the specificity of the exercise stimulus. Most of the resistance training studies in older adults involve single-joint exercises and the use of resistance machines, which limit the training movement to a fixed pattern in a single plane of motion. Although this regimen is effective at enhancing maximal muscular strength, it seems to elicit a more modest effect on functional performance (39,43). Our training intervention involved resistance training exercises that mimic the biomechanical movement patterns of everyday life activities, such as rising from a chair (e.g., squat), climbing the stairs (e.g., split squat), and twisting to pick an item up off the floor (e.g., core rotation).

Furthermore, many movement deficits develop during later adulthood such as a lack of ankle ROM and reduced hip extension, which result in adverse gait kinematics and a decline in functionality (34). Training programs specifically targeting these age-related movement deficits have been shown to enhance gait velocity and center of mass kinematics in the STS transition (15,44). Favorable changes in walking speed and STS kinematics may aid in the performance of tasks such as the TUG. Therefore, the inclusion of specific mobility exercises in our intervention (designed primarily to increase ankle, hip, and thoracic spine ROM) might have contributed to the large improvements in functional performance. Further research is required to confirm the mechanistic changes that underpin improvements in functional tasks after training.

It is important that resistance training evokes changes that are clinically meaningful for the intended population. Changes in laboratory-based measurements after a resistance training intervention are designed to reflect changes in clinically meaningful endpoints (28). Because laboratory measurements are not clinically meaningful endpoints, they must be correlated with those that are to be considered valid (28). For example, performance in the SCT is associated with self-reported functional abilities in older adults (5) and the test involves the same movement patterns as climbing the stairs in a real-life setting. Although correlations cannot establish cause and effect, it is likely that the \sim 13% improvement in SCT performance would have a direct influence on an aging person's ability to climb a flight of stairs in day-today life. This magnitude of change is also consistent with other resistance training studies (~ 9 to 14%) (29,30,35) and exceeds the MDC_{95%} recorded previously (7.7%) (40), confirming that the change was not due to measurement error or variation within individual performance. Further work is warranted to delineate a causal relationship between improvements in laboratory-based measurements and changes in day-to-day function.

The resistance training program followed the principle of progressive overload by systematically increasing resistance (grade of elastic band) and volume (number of sets and/or repetitions) over time. In addition, in accordance with NSCA guidelines (27), the difficulty of exercise selection was individually tailored according to the participant's ability and perception of effort. That is, exercises were modified using exercises of similar movement patterns but with different technical difficulties. For example, the body weight squat was progressed to a body weight lunge when the participant rated the squat exercise as "easy" (≤ 3 on the modified RPE Scale). Both movements are multijointed motor actions involving large muscle groups, but the lunge is unilateral in nature, reduces the base of support from a biped stance to a split stance, and requires greater hip flexor ROM. The lunge also necessitates a larger amount of muscle force to decelerate the body's inertia and then accelerate the body back to the starting position. Advancing from low-skill to high-skill exercises may improve movement quality to a greater extent than increasing resistance load or volume alone. Indeed, a ceiling effect exists whereby further increases in strength will not lead to additional functional improvements in older adults (6). Although modifying exercise selection based on individual ability is common practice in athlete populations, it is a strategy seldom included within training interventions for older adults. Researchers and practitioners should consider focusing on the primary movement pattern rather than the exercise itself, and move away from prescribing homogenous training programs for a largely heterogeneous population.

The increases in functional ability and IMTP strength were similar between SUP and UNSUP groups. Between-group comparisons did not reach statistical significance for any variable, which is further supported by the 95% CIs spanning zero. Although these nonsignificant results do not establish equivalence, the data imply that equivalency cannot be ruled out. This finding is in contrast to a recent meta-analytic review suggesting that, in a pooled analysis of 5 studies, SUP resistance training improves proxies of muscle strength to a greater extent than UNSUP programs (SMD = 0.51) (37). However, when considering the primary data, 3 of the 5 studies included in the review reported no differences between SUP and UNSUP interventions (1,19,46). Another included study compared SUP high-intensity training vs. a UNSUP low-intensity program (50); consequently, the difference between groups may be attributed to different loading strategies rather than the level of supervision. The remaining study reported larger improvements in function after 12 weeks of SUP strength and balance training compared with a parallel home-based program (36). Therefore, despite the recent publication of a well-designed meta-analysis (37), existing research comparing SUP vs. UNSUP resistance training programs in aging adults remains equivocal. Our data suggest that home-based resistance training with telephone support is an effective alternative to SUP programs, although this finding requires replication in interventions lasting several months rather than weeks.

This study is the first to demonstrate greater elevations in heart rate when untrained aging adults receive supervision during a resistance training intervention. Weekly telephone calls to the UNSUP group revealed lower mean stages of exercise progression compared with the SUP group, which implies that the greater heart rates may have been related to the completion of more advanced exercises. Direct supervision may have also fostered a higher quality in the execution of exercises due to continual technical feedback. Alternatively, the greater heart rates may be related to psychological factors such as competitiveness (i.e., presence of an audience) or external motivation (i.e., real-time encouragement). Interestingly, the average heart rate elicited in the SUP group (117 \pm 8 b·min⁻¹) was equivalent to ~70% of agepredicted HRmax (220-age), which meets the American College of Sports Medicine guidelines for moderateintensity aerobic exercise (24). The capacity of resistance training to contribute to the aerobic component of international physical activity guidelines has been reported recently (9) and suggests that this resistance training, when programed appropriately, can provide stimuli for both cardiovascular and musculoskeletal adaptation. In light of an increasingly sedentary population, promoting resistance training as a single method to achieve discernible health benefits should be considered. Future research should evaluate whether the higher heart rates elicited in the SUP vs. UNSUP group translate into greater improvements in cardiovascular fitness.

After exercise cessation, training-derived improvements were robust and remained above baseline values in both intervention groups. For example, performance in the 30-second chair STS test after DET was still $\sim 14\%$ greater than baseline. Previous studies have also reported that, after DET phases of 6–12 weeks, STS performance remains ~ 10 to 22% greater than pretraining values (2,14,36). Less retention of 30-second chair STS performance ($\sim 8\%$) has been

observed after longer DET periods of 24 weeks (48) and 1 year (18). It is likely that the residual effect of resistance training diminishes with longer periods of DET. Age may also mediate the effects of DET; Seco et al. (45) have previously reported better maintenance of balance performance among 65-74-year-olds compared with those aged 75 years or older. Given that we included younger participants (53.6 years) than the aforementioned studies (~ 65 years) (18,48), it might be expected that our subjects would retain a greater proportion of their training improvements. By contrast, the initial training regimen does not seem to influence DET's effect on functional performance. We found the residual benefit of resistance training was similar between SUP and UNSUP interventions, which is consistent with data obtained recently by Lacroix et al. (36). Others have also demonstrated that DET is not affected by training load, training duration, or repetition velocity among older adults (31,48). However, comparing posttraining to post-DET, the significant decreases in some parameters of physical performance highlight the negative effects of discontinuing a resistance training program. This reinforces the notion that aging individuals should be engaged in a regimen of resistance training across the lifespan to mitigate age-related declines in function.

A limitation of this study is that the investigator was not blinded to group allocation, although all participants received the same instructions and strictly adhered to a predetermined testing protocol. In addition, training intensity was controlled indirectly by selecting a target number of repetitions associated with a subjective perception of effort. Although resistance training load is usually quantified using a percentage of 1 repetition maximum, the use of RPE has been shown to be a valid indicator of elastic resistance training intensity in older adults (17). The weekly telephone support provided to the UNSUP group may also have encouraged exercise adherence (8). Therefore, it is unknown whether the same results would have occurred if there was no contact with participants during the intervention period. Furthermore, we did not include an inactive control group, although we have interpreted the magnitude of effects in relation to the error of measurements that were matched for time in our laboratory (4 weeks separating trials) (40). Finally, participants in this study were healthy adults aged 53.6 \pm 3.6 years (range: 50-62 years) and may not be representative of all elderly persons. Most previous studies have included adults aged above 65 years; so, comparisons made between our results and the current body of literature should take this age difference into consideration. The hypertrophic response to resistance training may be diminished with advancing age, but aging does not seem to impair one's ability to increase muscle strength (49). Future studies should assess whether a functional resistance training program with minimal supervision is well-tolerated by older and mobility-limited individuals.

To conclude, this study demonstrated that a 4-week functional resistance training program, performed using body

weight movements and elastic bands, elicited meaningful improvements in physical performance. The increases in functional ability and muscle strength were similar between SUP and UNSUP groups, suggesting that home-based resistance training is a practical and effective alternative to SUP programs for aging adults. Importantly, the training-induced improvements were largely preserved after exercise cessation.

PRACTICAL APPLICATIONS

A functional resistance training program may be implemented into clinical practice to mitigate age-related declines in muscle strength and function. Owing to the comparative effectiveness of SUP and UNSUP groups, our data also suggest that practitioners may prescribe home-based resistance training as a cost-effective and practical alternative to SUP programs that may help circumvent many barriers to physical activity in the aging population, such as lack of time, money, and transportation. This finding, however, requires replication in interventions lasting several months rather than weeks. The adaptations to a resistance training program are well maintained beyond the cessation of training, although lifelong participation in resistance training should be encouraged to attenuate the inevitable decline in functional capacity during later adulthood. Taken together, these findings suggest that aging adults should choose a preferable environment for exercise (i.e., UNSUP at home or SUP in a facility) that will foster consistent adherence to resistance training in the longer term.

ACKNOWLEDGMENTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. All data were collected in the Sport, Health and Exercise Science laboratory at the University of Hull.

REFERENCES

- Almeida, TL, Alexander, NB, Nyquist, LV, Montagnini, ML, Santos, ACS, Rodrigues, GHP, Negrão, CE, Trombetta, IC, and Wajngarten, M. Minimally supervised multimodal exercise to reduce falls risk in economically and educationally disadvantaged older adults. *J Aging Phys Act* 21: 241–259, 2013.
- Ansai, JH, Aurichio, TR, Gonçalves, R, and Rebelatto, JR. Effects of two physical exercise protocols on physical performance related to falls in the oldest old: A randomized controlled trial. *Geriatr Gerontol Int* 16: 492–499, 2016.
- Ashworth, NL, Chad, KE, Harrison, EL, Reeder, BA, and Marshall, SC. Home versus center based physical activity programs in older adults. *Cochrane Database Syst Rev* 1: CD004017, 2005.
- ATS. ATS statement: Guidelines for the six-minute walk test. Am J Respir Crit Care Med 166: 111–117, 2002.
- Baldwin, JN, McKay, MJ, Hiller, CE, Moloney, N, Nightingale, EJ, and Burns, J. Relationship between physical performance and selfreported function in healthy individuals across the lifespan. *Musculoskelet Sci Pract* 30: 10–17, 2017.
- Bean, JF, Vora, A, and Frontera, WR. Benefits of exercise for community-dwelling older adults. *Arch Phys Med Rehabil* 85(7 Suppl 3): S31–S42, 2004.

- Benavent-Caballer, V, Sendín-Magdalena, A, Lisón, JF, Rosado-Calatayud, P, Amer-Cuenca, JJ, Salvador-Coloma, P, and Segura-Ortí, E. Physical factors underlying the Timed "Up and Go" test in older adults. *Geriatr Nurs* 37: 122–127, 2016.
- Bennell, KL, Campbell, PK, Egerton, T, Metcalf, B, Kasza, J, Forbes, A, Bills, C, Gale, J, Harris, A, Kolt, GS, Bunker, SJ, Hunter, DJ, Brand, CA, and Hinman, RS. Telephone coaching to enhance a home-based physical activity program for knee osteoarthritis: A randomized clinical trial. *Arthritis Care Res (Hoboken)* 69: 84–94, 2017.
- Bharti, N, Hrubeniuk, T, Mayo, A, Senechal, M, and Bouchard, DR. Resistance training contribute to the aerobic components of an exercise session in adults but not as much in older adults. *Int J Exerc Sci* 10: 406–416, 2017.
- Bozic, PR, Pazin, NR, B erjan, BB, Planic, NM, and Cuk, ID. Evaluation of the field tests of flexibility of the lower extremity: Reliability and the concurrent and factorial validity. *J Strength Cond Res* 24: 2523–2531, 2010.
- Byrne, C, Faure, C, Keene, DJ, and Lamb, SE. Ageing, muscle power and physical function: A systematic review and implications for pragmatic training interventions. *Sports Med* 46: 1311–1332, 2016.
- Cadenas-Sanchez, C, Sanchez-Delgado, G, Martinez-Tellez, B, Mora-Gonzalez, J, Löf, M, España-Romero, V, Ruiz, JR, and Ortega, FB. Reliability and validity of different models of TKK hand dynamometers. *Am J Occup Ther* 70: 7004300010, 2016.
- Carli, F and Scheede-Bergdahl, C. Prehabilitation to enhance perioperative care. *Anesthesiol Clin* 33: 17–33, 2015.
- Carvalho, MJ, Marques, E, and Mota, J. Training and detraining effects on functional fitness after a multicomponent training in older women. *Gerontology* 55: 41–48, 2009.
- Choi, JH and Kim, NJ. The effects of balance training and ankle training on the gait of elderly people who have fallen. *J Phys Ther Sci* 27: 139–142, 2015.
- 16. Coetsee, C and Terblanche, E. The time course of changes induced by resistance training and detraining on muscular and physical function in older adults. *Eur Rev Aging Phys Act* 12: 7, 2015.
- Colado, JC and Triplett, NT. Effects of a short-term resistance program using elastic bands versus weight machines for sedentary middle-aged women. J Strength Cond Res 22: 1441–1448, 2008.
- Correa, CS, Cunha, G, Marques, N, Oliveira-Reischak, Ã, and Pinto, R. Effects of strength training, detraining and retraining in muscle strength, hypertrophy and functional tasks in older female adults. *Clin Physiol Funct Imaging* 36: 306–310, 2015.
- Cyarto, EV, Brown, WJ, Marshall, AL, and Trost, SG. Comparison of the effects of a home-based and group-based resistance training program on functional ability in older adults. *Am J Health Promot* 23: 13–17, 2008.
- Devereux-Fitzgerald, A, Powell, R, Dewhurst, A, and French, DP. The acceptability of physical activity interventions to older adults: A systematic review and meta-synthesis. *Soc Sci Med* 158: 14–23, 2016.
- Dias, CP, Toscan, R, de Camargo, M, Pereira, EP, Griebler, N, Baroni, BM, and Tiggemann, CL. Effects of eccentric-focused and conventional resistance training on strength and functional capacity of older adults. *Age* 37: 99, 2015.
- Dos'Santos, T, Thomas, C, Jones, PA, McMahon, JJ, and Comfort, P. The effect of hip joint angle on isometric mid-thigh pull kinetics. J Strength Cond Res 31: 2748–2757, 2017.
- Franco, MR, Howard, K, Sherrington, C, Ferreira, PH, Rose, J, Gomes, JL, and Ferreira, ML. Eliciting older people's preferences for exercise programs: A best-worst scaling choice experiment. J Physiother 61: 34–41, 2015.
- 24. Garber, CE, Blissmer, B, Deschenes, MR, Franklin, BA, Lamonte, MJ, Lee, IM, Nieman, DC, and Swain, DP. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc* 43: 1334–1359, 2011.

VOLUME 33 | NUMBER 10 | OCTOBER 2019 | 2741

- Glenn, JM, Gray, M, and Binns, A. The effects of loaded and unloaded high-velocity resistance training on functional fitness among community-dwelling older adults. *Age Ageing* 44: 926–931, 2015.
- 26. Gruther, W, Wick, F, Paul, B, Leitner, C, Posch, M, Matzner, M, Crevenna, R, and Ebenbichler, G. Diagnostic accuracy and reliability of muscle strength and endurance measurements in patients with chronic low back pain. *J Rehabil Med* 41: 613–619, 2009.
- Haff, GG and Triplett, TN. Program design for resistance training. In: *Essentials of Strength Training and Conditioning*. Champaign, IL: Human Kinetics, 2015. pp. 439–469.
- Halperin, I, Vigotsky, AD, Foster, C, and Pyne, DB. Strengthening the practice of exercise and sport science. *Int J Sports Physiol Perform* 13: 127–134, 2018.
- 29. Henwood, TR and Taaffe, DR. Short-term resistance training and the older adult: The effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging* 26: 305–313, 2006.
- Henwood, TR, Riek, S, and Taaffe, DR. Strength versus muscle power-specific resistance training in community-dwelling older adults. J Gerontol A Biol Sci Med Sci 63: 83–91, 2008.
- Henwood, TR and Taaffe, DR. Detraining and retraining in older adults following long-term muscle power or muscle strength specific training. J Gerontol A Biol Sci Med Sci 63: 751–758, 2008.
- Hintermeister, RA, Lange, GW, Schultheis, JM, Bey, MJ, and Hawkins, RJ. Electromyographic activity and applied load during shoulder rehabilitation exercises using elastic resistance. *Am J Sports Med* 26: 210–220, 1998.
- Jones, CJ, Rikli, RE, and Beam, WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport* 70: 113–119, 1999.
- Jung, H and Yamasaki, M. Association of lower extremity range of motion and muscle strength with physical performance of community-dwelling older women. J Physiol Anthropol 35: 30, 2016.
- Kalapotharakos, VI, Michalopoulos, M, Tokmakidis, SP, Godolias, G, and Gourgoulis, V. Effects of a heavy and a moderate resistance training on functional performance in older adults. *J Strength Cond Res* 19: 652–657, 2005.
- 36. Lacroix, A, Kressig, RW, Muehlbauer, T, Gschwind, YJ, Pfenninger, B, Bruegger, O, and Granacher, U. Effects of a supervised versus an unsupervised combined balance and strength training program on balance and muscle power in healthy older adults: A randomized controlled trial. *Gerontology* 62: 275–288, 2016.
- Lacroix, A, Hortobagyi, T, Beurskens, R, and Granacher, U. Effects of supervised vs. unsupervised training programs on balance and muscle strength in older adults: A systematic review and metaanalysis. *Sports Med* 47:2341–2361, 2017.
- Liu, CJ and Latham, NK. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev* 3: Cd002759, 2009.

- Liu, CJ and Latham, N. Can progressive resistance strength training reduce physical disability in older adults? A meta-analysis study. *Disabil Rehabil* 33: 87–97, 2011.
- Northgraves, MJ, Hayes, SC, Marshall, P, Madden, LA, and Vince, RV. The test-retest reliability of four functional mobility tests in apparently healthy adults. *Isokinet Exerc Sci* 24: 171–179, 2016.
- Orange, S and Smith, A. Evidence-based strength and conditioning in soccer. *Health Fitness J Can* 9: 21–37, 2016.
- Papa, EV, Dong, X, and Hassan, M. Resistance training for activity limitations in older adults with skeletal muscle function deficits: A systematic review. *Clin Interv Aging* 12: 955–961, 2017.
- Peterson, MD, Rhea, MR, Sen, A, and Gordon, PM. Resistance exercise for muscular strength in older adults: A meta-analysis. *Ageing Res Rev* 9: 226–237, 2010.
- Schot, PK, Knutzen, KM, Poole, SM, and Mrotek, LA. Sit-to-stand performance of older adults following strength training. *Res Q Exerc Sport* 74: 1–8, 2003.
- 45. Seco, J, Abecia, LC, Echevarria, E, Barbero, I, Torres-Unda, J, Rodriguez, V, and Calvo, JI. A long-term physical activity training program increases strength and flexibility, and improves balance in older adults. *Rehabil Nurs* 38: 37–47, 2013.
- 46. Van Roie, E, Delecluse, C, Opdenacker, J, De Bock, K, Kennis, E, and Boen, F. Effectiveness of a lifestyle physical activity versus a structured exercise intervention in older adults. *J Aging Phys Act* 18: 335–352, 2010.
- 47. Van Roie, E, Delecluse, C, Coudyzer, W, Boonen, S, and Bautmans, I. Strength training at high versus low external resistance in older adults: Effects on muscle volume, muscle strength, and forcevelocity characteristics. *Exp Gerontol* 48: 1351–1361, 2013.
- Van Roie, E, Walker, S, Van Driessche, S, Baggen, R, Coudyzer, W, Bautmans, I, and Delecluse, C. Training load does not affect detraining's effect on muscle volume, muscle strength and functional capacity among older adults. *Exp Gerontol* 98: 30–37, 2017.
- Walker, S and Häkkinen, K. Similar increases in strength after short-term resistance training due to different neuromuscular adaptations in young and older men. J Strength Cond Res 28: 3041–3048, 2014.
- Watson, SL, Weeks, BK, Weis, LJ, Horan, SA, and Beck, BR. Heavy resistance training is safe and improves bone, function, and stature in postmenopausal women with low to very low bone mass: Novel early findings from the LIFTMOR trial. *Osteoporos Int* 26: 2889– 2894, 2015.
- 51. Wright, AA, Cook, CE, Baxter, GD, Dockerty, JD, and Abbott, JH. A comparison of 3 methodological approaches to defining major clinically important improvement of 4 performance measures in patients with hip osteoarthritis. *J Orthop Sports Phys Ther* 41: 319– 327, 2011.

2742 Journal of Strength and Conditioning Research